

Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics



CrossMark
click for updates

www.renhyd.org



RESEARCH ARTICLE

Sleep Quality and Associated Factors among Adolescents

Asude Rabia Ozkan^{a,*}, Ozge Kucukerdonmez^a, Gulsah Kaner^b

^aDepartment of Nutrition and Dietetics, Institute of Health Sciences, Ege University, İzmir, Turkey.

^bDepartment of Nutrition and Dietetics, Faculty of Health Sciences, İzmir Katip Çelebi University, İzmir, Turkey.

*asuderabia@hotmail.com

Assigned Editor: Tania Fernández Villa. Universidad de León. España.

Received: 08/03/2020; accepted: 01/06/2020; published: 03/07/2020.

KEYWORDS

Adolescent;
Feeding Behavior;
Sleep;
Sleep Hygiene;
Anthropometry.

➤ Sleep Quality and Associated Factors among Adolescents

ABSTRACT

Introduction: This study aimed to investigate sleep quality and sleep patterns in adolescents. The second aim was to find out whether there was an association between sleep quality, sleep patterns, dietary habits, food consumptions, and anthropometric measurements in an adolescent population.

Material and Methods: This cross-sectional study was conducted on 346 adolescents. Data was recorded with a questionnaire form. The questionnaire form included items on adolescents' characteristics, sleep patterns, dietary habits, food consumptions, and anthropometric measurements. Sleep quality was measured using the Pittsburgh Sleep Quality Index (PSQI).

Results: Sleep duration of >8 hours (67.6%) and sleep latency of 15 minutes (53.8%) were the most commonly identified sleeping patterns. The mean PSQI score was 3.07±2.54. PSQI scores revealed poor sleep quality in 13.6% of participants. A significant difference was observed between age, disease diagnosed by physician, skipping main meals, having regular breakfast, and PSQI score ($p<0.05$). PSQI score was found to be significantly associated with spending time in front of the computer and regular sleep. The amount of daily saturated fatty acid was statistically significantly lower ($p=0.040$) in individuals in the good PSQI.

Conclusions: The vast majority of adolescents had good sleep quality. Sleep duration of adolescents was consistent with the recommended need. Individual factors, dietary habits, food consumption, and screen times were factors associated with sleep quality. This current study results support the development of interventions to help adolescents improve sleep quality. We recommend further investigation to clarify this finding.

Calidad del sueño y factores asociados entre los adolescentes

PLABRAS CLAVE

Adolescente;
Conducta Alimentaria;
Sueño;
Higiene del Sueño;
Antropometría.

RESUMEN

Introducción: Este estudio tuvo como objetivo investigar la calidad del sueño y los patrones de sueño en los adolescentes. El segundo objetivo fue averiguar si había una asociación entre la calidad del sueño, los patrones de sueño, los hábitos alimenticios, el consumo de alimentos y las mediciones antropométricas en una población adolescente.

Material y Métodos: Este estudio transversal se realizó en 346 adolescentes. Los datos se registraron con un formulario de cuestionario. El formulario del cuestionario incluía elementos sobre las características de los adolescentes, los patrones de sueño, los hábitos alimenticios, el consumo de alimentos y las mediciones antropométricas. La calidad del sueño se midió utilizando el Índice de calidad del sueño de Pittsburgh (PSQI).

Resultados: La duración del sueño >8 horas (67,6%) y la latencia del sueño de 15 minutos (53,8%) fueron los patrones de sueño más comúnmente identificados. La puntuación media de PSQI fue de $3,07 \pm 2,54$. Las puntuaciones del PSQI revelaron una mala calidad del sueño en el 13,6% de los participantes. Se observó una diferencia significativa entre la edad, la enfermedad diagnosticada por el médico, la omisión de las comidas principales, el desayuno regular y el puntaje PSQI ($p < 0,05$). Se encontró que el puntaje del PSQI estaba significativamente asociado con pasar tiempo frente a la computadora y dormir regularmente. La cantidad de ácido graso saturado diario fue menor estadísticamente significativa ($p = 0,040$) en individuos con buen PSQI.

Conclusiones: La gran mayoría de los adolescentes tenían buena calidad del sueño. La duración del sueño de los adolescentes fue consistente con la necesidad recomendada. Los factores individuales, los hábitos alimenticios, el consumo de alimentos y los tiempos de pantalla fueron factores asociados con la calidad del sueño. Los resultados de este estudio actual apoyan el desarrollo de intervenciones para ayudar a los adolescentes a mejorar la calidad del sueño. Recomendamos una mayor investigación para aclarar este hallazgo.

CITATION

Ozkan AR, Kucukerdonmez O, Kaner G. Sleep Quality and Associated Factors among Adolescents. Rev Esp Nutr Hum Diet. 2020; 24(3): 256-67. doi: 10.14306/renhyd.24.3.1017

INTRODUCTION

A regular sleep is one of the basic elements of a healthy life for both children and adults. Many somatic, cognitive and psychological processes are strongly influenced by good sleep, and good sleep contributes to improving health¹. It has been shown in many meta-analysis that chronic insufficient sleep (<6 hours) has negative effects on metabolism and it increases the risk of type 2 diabetes, obesity, hypertension, cardiovascular disease and mortality²⁻⁷. Despite the fact that adolescents need 8 to 10 hours of sleep⁸, the National Sleep Foundation has reported that they tend to have irregular sleep patterns across the week and only the 15% of them have an

optimal sleep duration⁹. Sleep has an important role in brain development, and inadequate amount of sleep in adolescence may affect hypothalamic mechanisms that regulate appetite and energy metabolism¹⁰.

Studies have reported that short sleep duration is associated with the risk of obesity^{11,12}. Adequate sleep duration is critical for preventing obesity. Overweight and obesity in adolescents not only increase the risk of chronic diseases and psychosocial problems such as decreased self-esteem, poor body image, and social exclusion, but also neuropsychological dysfunctions such as depression¹³. Furthermore, short sleep duration may have an impact on food intake and appetite. Changes in lifestyle and unhealthy habits such as following a high-calorie diet

are often associated with altered sleep patterns and sleep efficiency¹⁴. In addition to this, it has been generally demonstrated that the habit of snacking is related to the shortness of sleep duration; nevertheless, it is not clear whether this is due to frequent eating during the day or the high energy and low nutrient content of snacks. However, it is certain that there is a positive relationship between short and irregular sleep patterns and an unhealthy diet¹⁵.

Daily activities, individual factors, and environmental changes can have an impact on the sleep patterns of adolescents. Since sleep duration and quality are of great importance in adolescents, the aim of this study was to determine sleep quality and sleep patterns among 11-13 years old adolescents. The second aim of the present study was to find out whether there was an association between sleep quality, sleep patterns, dietary habits, food consumptions, and anthropometric measurements in an adolescent population.

MATERIAL AND METHODS

Study design

This study conducted on adolescents between the ages of 11-13 from two public schools between September and December 2017. For this research, approval from Ethics Committee of İzmir Katip Çelebi University were obtained (Decision No. 22.06.2017-95). All participants gave their assent in writing and necessary permissions were obtained from their parents.

Participants and sample

There were 3484 students in the city center of Muğla. A cross-sectional study was carried out in a random sample of adolescents who were attending two public schools in Muğla. The participants were selected using a stratified proportional sampling method according to the number of sampling calculated by the known population with a 95% confidence interval. The sample group included 346 participants. Within this group, the 11-year age group consisted of 65 girls and 70 boys; the age group of 12 consisted of 51 girls, 53 boys; and the age group of 13 consisted of 56 girls and 53 boys.

Questionnaire form

Data was recorded with a questionnaire form applied via face-to face method. The questionnaire form included

items on adolescents' characteristics (age, gender, disease diagnosed by physician), screen times (spending time in front of computer and television in leisure time) dietary habits (regular breakfast, skipping main meals), sleep patterns (sleep duration, sleep latency, habitual sleep efficiency), sleep quality, anthropometric measurements, and food consumptions.

Sleep quality

Pittsburgh sleep quality index (PSQI) was applied to determine the sleep quality. In the form consisting of 18 questions, each question is evaluated with a number from 0 to 3. The sum of the scores gives the total PSQI score. The sleep quality of those with less than 5 points in total is "good" whereas that of those 5 points or above is considered as "poor"^{16,17}.

Anthropometric measurements

The body weight, height, waist circumference, and hip circumference were evaluated by previously standardized nutritionists who used conventional anthropometric techniques. In addition, their waist/hip and waist/height ratios were calculated. A portable scale was used to measure body weight. Height measurement was performed using a 2m long inflexible steel waist circumference was measured with a flexible tape¹⁸. The anthropometric measurements were assessed using the WHO Growth Reference for 5-19 Years-2007¹⁹. Gender and age specific z-scores were calculated using the WHO AnthroPlus software²⁰. The children were classified into five categories of BMI for age Z-score (BAZ): underweight, at risk of underweight, normal weight, overweight and obese, in accordance with the cut-off points of < (-2SD), (-2SD) to (-1SD), (-1SD) to (1SD), 1SD, 1SD to 2SD and ≥2SD Z-scores, respectively¹⁹.

Food consumptions

The dietary intake was evaluated based on 24-hour food recall. The 24-hour recall was undertaken in chronological order of consumption (from morning to night). Participants were asked to record all the foods and beverages during the previous day. The portion contents of the meals consumed were calculated by using the book called Standard Recipes for Institutions²¹. The number of grams of the foods specified in the records as a standard was calculated by using the book called "Foods and Meals Photo Catalog: Standards and Quantities"²². Daily food consumption was noted, and daily energy, micro and macronutrient intakes were identified by the Nutrition Information System Software (BEBİS)²³.

Statistical analysis

All analyses were done using Statistical Package for the Social Sciences version 24.0 (SPSS Inc. Chicago, IL, USA)²⁴. Frequency tables and descriptive statistics were used to interpret the results. Following the parametric methods, T test was used to compare the independent groups. Following the non-parametric methods, the Mann-Whitney U test was used to compare the two independent groups, and the Kruskal-Wallis H test was used to compare the three or more independent groups.

RESULTS

Adolescent's characteristics

Of the 346 adolescents included in the study, 50.9% were boys, 49.1% were girls, and the mean age was 11.9±0.8 years. It was determined that 86.1% of the individuals did not have a disease diagnosed by the physician. The most common diseases diagnosed by the physician were respiratory system diseases (25.0%) and cardiovascular diseases (18.8%) (Table 1).

Screen times

Spending 2-3 hours in front of computer and television in leisure time was 21.9% and 36.1%, respectively (Table 1).

Table 1. Adolescent characteristics and sleep patterns (n=346).

Age (year), mean (SD)	11.9 (0.8)
Age group (year), n (%)	
11	135 (39.0)
12	104 (30.0)
13	107 (30.9)
Gender, n (%)	
Female	170 (49.1)
Male	176 (50.9)
Disease diagnosed by physician, n (%)	
Yes	48 (13.9)
No	298 (86.1)

Diseases, n (%)	
Cardiovascular diseases	9 (18.8)
Diabetes	3 (6.3)
Mental health problems	7 (14.6)
Respiratory diseases	12 (25.0)
Muscular system problems	3 (6.3)
Endocrine diseases	7 (14.5)
Vitamin-mineral deficiencies	7 (14.5)
Spending time in front of computer	
Never	92 (26.7)
< 1	88 (25.4)
1	63 (18.2)
2-3	76 (21.9)
> 3	27 (7.8)
Spending time in front of television	
Never	27 (7.8)
< 1	64 (18.5)
1	97 (28.0)
2-3	125 (36.1)
> 3	33 (9.6)
Skipping main meal	
Yes	35 (10.1%)
No	217 (62.8%)
Sometimes	94 (27.1%)
Regular breakfast	
Yes	203 (58.7%)
No	31 (9.0%)
Sometimes	112 (32.3%)
Sleep duration (hour), n (%)	
3-5	11 (3.2)
6-8	101 (29.2)
> 8	234 (67.6)
Sleep latency, n (%)	
0-15 minimum	186 (53.8)
15-30 minimum	107 (30.9)
31-60 minimum	35 (10.1)
> 60 minimum	18 (5.2)
Habitual sleep efficiency, n (%)^a	
≥ 85%	334 (96.5)
75-84%	6 (1.7)
65-74%	4 (1.2)
< 65%	2 (0.6)
PSQI total score, n (%)	
Good (< 5)	299 (86.4)
Poor (≥ 5)	47 (13.6)

SD: Standard deviation:
a: (total of hours asleep)/(total of hours in bed) x100.

Dietary habits

More than half (58.7%) of the adolescents eat breakfast regularly and 27.1% of them declared that they sometimes skipped main meals (Table 1).

Sleep patterns and sleep quality

Sleep duration of >8 hours (67.6%), sleep latency of 15 minutes (53.8%) were the most commonly identified sleeping patterns. The mean PSQI score was 3.07±2.54. PSQI scores revealed poor sleep quality in 13.6% of participants. Habitual sleep efficiency was ≥85% in 96.5% of subjects (Table 1).

Anthropometric measurements

The mean of body weight, height, waist circumference, hip circumference, waist to hip and waist to height ratio of adolescents were 46.2±11.7kg, 153.8±8.5cm, 69.4±9.6cm, 84.0±9.4cm, 0.8±0.0, 0.4±0.0, respectively. In the evaluation of BAZ, the prevalence of overweight and obesity was found to be 10.4% and 8.4%, respectively (Table 2).

group of 11. Moreover, individuals in the age group of 13 had poorer sleep quality compared to the age group of 12. Individuals with a disease had poorer sleep quality than those without a disease (p<0.05).

Screen times: A statistically significant difference was found between those who spent 2-3 hours, those who spent 1 or less than 1 hour, and those who spent no time in front of a computer, in terms of PSQI score (p=0.001).

Dietary habits: Individuals who did not have regular breakfast had poorer sleep quality compared to those who regularly and sometimes had breakfast. Those who skipped meals had poorer sleep quality than those who did not. It was determined that individuals who slept 3-5 hours a day had poorer sleep quality compared to those who slept more than 6-8 and 8 hours (p<0.05) (Table 4).

Anthropometric measurements: There was no statistically significant relationship between PSQI score and body weight, BMI classification, waist circumference, hip circumference, and waist/height ratio (p>0.05) (Table 5).

Table 2. Anthropometric measurements of adolescents.

	Boys (n=176) Mean±SD	Girls (n=170) Mean±SD	Total Mean±SD
Body weight (kg)	46.2±12.0	46.2±11.4	46.2±11.7
Height (cm)	153.4±8.9	154.2±8.0	153.8±8.5
WC (cm)	71.4±9.8	67.4±8.9	69.4±9.6
HC (cm)	83.1±9.2	84.9±9.4	84.0±9.4
WC/HC	0.8±0.0	0.7±0.0	0.8±0.0
WC/height	0.4±0.0	0.4±0.0	0.4±0.0

WC: Waist circumference; HC: Hip circumference; NC: Neck circumference.

Food consumptions

On average, adolescents consumed 1875.6 calories, with caloric intake comprised of 49.3% carbohydrates, 34.8% fat, and 15.8% protein. Intakes of energy, carbohydrate, protein, fat, vitamin A, riboflavin, niacin, vitamin B6, vitamin B12, sodium, zinc, and iron were higher in boys (p<0.05) (Table 3).

Factors Associated with Sleep Quality

Adolescent’s characteristics: Individuals in the age group of 13 and 12 had poorer sleep quality compared to the age

There was a weak negative relationship between PSQI and the sleep duration (r=0.499; p=0.000), a weak positive relationship between PSQI and height (r=0.185; p=0.001), and a weak negative relationship between PSQI and waist/hip ratio (r=-0.164; p=0.002) (Table 6).

Food consumptions: When the daily energy and nutrient intake values of individuals were compared to PSQI, only the amount of saturated fatty acid taken daily was statistically significantly lower (p=0.040) in individuals in the good PSQI class compared to those in the poor PSQI class (Table 7).

Table 3. Daily energy and nutrient intakes of adolescents.

	Boys (n=176) Mean±SD	Girls (n=170) Mean±SD	Total Mean±SD	P
Energy (kcal/day)	1956.8±228.6	1791.1±276.9	1875.6±266.3	<0.001
Carbohydrate (%)	50.3±6.9	48.2±7.9	49.3±7.5	0.010
Protein (%)	16.6±4.0	15.0±3.3	15.8±3.8	<0.001
Fat (%)	33.0±5.4	36.6±7.3	34.8±6.6	<0.001
SFA (g)	25.9±8.0	27.5±10.1	26.7±9.1	0.362
Fiber (g)	19.9±6.6	20.6±7.4	20.3±7.0	0.414
Vitamin A (µg)	732.2±550.5	917.6±346.3	898.9±897.3	<0.001
Vitamin E (mg)	13.1±5.6	10.20±2.9	13.2±5.7	0.977
Vitamin C (mg)	99.6±84.0	10.8±2.5	104.8±80.8	0.116
Thiamine (mg)	0.8±0.2	917.6±346.3	0.8±0.2	0.446
Riboflavin (mg)	1.5±0.5	1.2±2.9	1.4±0.4	0.011
Niacin (mg)	11.0±5.0	10.8±2.5	10.1±4.5	<0.001
Vitamin B6 (mg)	1.2±0.3	1.1±0.3	1.2±0.3	0.010
Folate (µg)	114.7±33.6	115.1±34.4	114.9±34.0	0.903
Vitamin B12 (µg)	5.2±8.3	3.6±2.3	4.4±6.1	0.001
Sodium (mg)	4408.1±1070.2	3783.3±1589.0	4101.1±1384.1	<0.001
Potassium (mg)	2305.0±588.0	2331.5±650.4	2318.0±618.7	0.558
Magnesium (mg)	253.1±58.0	249.9±65.2	251.6±61.6	0.428
Calcium (mg)	917.6±346.3	869.1±318.3	893.8±333.2	0.238
Zinc (mg)	10.2±2.9	8.9±2.7	9.5±2.9	<0.001
Iron (mg)	10.8±2.5	10.3±2.8	10.6±2.7	0.049

SFA: Saturated fatty acid.

DISCUSSION

The need for sleep may vary depending on individual and environmental factors such as age, gender, disease history, and lifestyle habits²⁵. In this study, 13.6% of adolescents had poor sleep quality. The poor sleep quality rate in adolescents was 20% in the Xu *et al.*'s study²⁶ and 54.7% in Şenol *et al.*'s study²⁷. Studies have reported that prolonged sleep latency and increased sleep problems in adolescents^{28,29}. Sleep latency was >30 minutes in 15.3% of adolescents in our cohort, which seem lower than previously reported rates^{28,29}. Habitual sleep efficiency was ≥85% in 96.5% of subjects in our cohort, which seems also higher than previously reported rates for ≥85%

habitual sleep efficiency in Turkish adolescents^{27,28}. The results are considered to be incompatible due to different measurement methods, different demographic, cultural characteristics and differences in the mean age. National Sleep Foundation states that adolescents should sleep 8-10 hours⁹. In our study, more than half of adolescents slept 8 hours or more. In our cohort, since the mean age of the students was 11.9±0.8 years, the daily sleep duration may be found to be 8 hours or more. The sleep duration detected was consistent with the recommendation of the National Sleep Foundation. However, it was determined in our cohort that there was a significant relationship between age and daily sleep duration, and that as the age increased, daily sleep duration decreased. Similarly, Felden *et al.* identified that the risk of poor sleep quality increased

Table 4. Comparison of some characteristics of individuals with PSQI score.

	n	Total PSQI		p
		Mean±SD	Median	
Gender				
Male	176	2.88±2.3	2.0	0.267
Female	170	3.28±2.7	3.0	
Age (year)				
11 ^(a)	135	2.30±2.1	2.0	<0.01 [a-b,c] [b-c]
12 ^(b)	104	3.29±2.3	2.5	
13 ^(c)	107	3.84±2.9	3.0	
BMI classification				
At risk of underweight	45	3.09±2.6	2.0	0.765
Normal weight	236	3.17±2.6	3.0	
Overweight	36	2.81±2.4	2.0	
Obese	29	2.57±1.6	2.0	
Disease diagnosed by physician				
Yes	48	4.08±3.2	3.5	0.023
No	298	2.92±2.3	2.0	
Skipping main meal				
Yes ^(a)	35	4.86±3.6	4.0	<0.01 [b-a,c]
No ^(b)	217	2.39±1.9	2.0	
Sometimes ^(c)	94	4.01±2.5	4.0	
Regular breakfast				
Yes ^(a)	203	2.54±1.9	2.0	<0.01 [b-a,c]
No ^(b)	31	5.19±3.7	4.0	
Sometimes ^(c)	112	3.47±2.6	3.0	
Spending time in front of computer in leisure time				
Never ^(a)	92	2.72±2.5	2.0	<0.01 [a,b,c-d]
< 1 ^(b)	88	2.45±1.8	2.0	
1 ^(c)	63	2.73±2.2	2.0	
2-3 ^(d)	76	4.22±2.9	4.0	
> 3 ^(e)	27	3.89±2.8	3.0	
Spending time in front of television in leisure time				
Never	27	3.81±2.7	3.0	0.200
< 1	64	2.94±2.7	2.0	
1	97	2.77±2.2	2.0	
2-3	125	3.09±2.6	2.0	
> 3	33	3.55±2.5	3.0	
Daily sleep duration (hour)				
3-5 ^(a)	11	9.82±2.7	10.0	<0.01 [a-b,c][b-c]
6-8 ^(b)	101	4.32±2.6	4.0	
> 8 ^(c)	234	2.22±1.6	2.0	

Mann-Whitney U test and Kruskal-Wallis H test were used.
Multiple significant differences between groups were shown by using the a,b,c,d,e characters.

Table 5. Comparison of anthropometric measurements of individuals by gender in terms of PSQI score.

	Boys (n=176)		P	Girls (n=170)		P
	Good PSQI (≤5)	Poor PSQI (>5)		Good PSQI (≤5)	Poor PSQI (>5)	
	Mean±SD	Mean±SD		Mean±SD	Mean±SD	
Body weight (kg)	46.4±12.3	44.6±9.7	0.582	45.6±11.3	49.4±11.7	0.089
Height (cm)	153.2±8.8	155.7±9.5	0.237	153.7±8.3	156.5±6.0	0.044
WC (cm)	71.7±10.1	69.1±7.1	0.330	67.1±8.9	68.6±8.8	0.328
HC (cm)	83.4±9.4	80.4±7.1	0.188	84.3±9.3	88.1±9.4	0.030
WC/HC	0.8±0.0	0.8±0.0	0.696	0.8±0.0	0.7±0.0	0.089
WC/height	0.4±0.0	0.4±0.0	0.097	0.4±0.0	0.4±0.0	0.664

WC: Waist circumference; HC: Hip circumference; NC: Neck circumference.
Mann-Whitney U test and Student's T test were used.

with age³⁰. This association may be explained by the fact that, while there is a cumulative increase of academic and recreational activities, there is also a reduction of time of sleep throughout the years³¹. Besides, the delay in the biological process of melatonin secretion is related with the advancement of puberty and with the reduced sleep hours among adolescents³².

This study has found that age does but gender does not affect sleep quality. Results of various studies report contradictory findings about whether boys or girls obtain more sleep. For example, one study found that girls obtained less sleep than boys and reported greater sleep disturbances³³. Another study showed that girls reported longer ideal sleep duration³⁴.

In the current study, a negative relation was detected between the amount of time the adolescents spent in front of a computer and sleep quality. Similar to our results, Akçay *et al.* showed that as the adolescents spent more time with their media products, their sleep quality deteriorated³⁵. The daily watch on the screen has increased significantly among adolescents. Excessive technology use may contribute to the adolescent insufficient sleep³⁶. A correlation was found between sleep deprivation and spending more than 2 hours in front of a TV or computer³⁷. Watching too much TV has been found to be associated with sleep disorders in adolescence, and the presence of a TV or a computer in the child's room has been shown to cause changes in sleep parameters³⁸. It has been emphasized by experts that children over the age of two should be limited to a maximum of 2 hours in front of the screen³⁹.

It has been identified that adolescents with good and bad sleep quality generally have high dietary fat and saturated fat intake, and low fiber intake in our study. However, no statistically significant difference was identified in terms of energy, macro and micronutrient intake, except for saturated fat intake. Studies have shown that short sleepers have higher energy intakes, notably from fat and snacks, than do normal sleepers^{40,41}. Insufficient sleep duration

Table 6. Correlation of PSQI scores with some parameters.

	PSQI	
	r	p
Sleep duration (hour)	-0.499	0.000
Energy (kcal)	-0.008	0.887
BMI for age Z score	0.015	0.776
Body weight (kg)	0.099	0.065
Height (cm)	0.185	0.001
WC (cm)	-0.003	0.957
HC (cm)	0.102	0.058
WC/HC	-0.164	0.002
WC/height	-0.086	0.109

WC: Waist circumference; HC: Hip circumference;
NC: Neck circumference.

*In cases where at least one of the two quantitative variables does not have distribution, Spearman correlation coefficient was used in evaluating the relationship between the variables

Table 7. Comparison of daily energy and nutrients in terms of PSQI classification.

	Good PSQI (≤ 5)		Poor PSQI (> 5)		p
	Mean \pm SD	Median	Mean \pm SD	Median	
Energy (kcal/day)	1876.6 \pm 268.2	1904.0	1867.8 \pm 257.0	1879.0	0.594
Carbohydrate (%)	49.3 \pm 7.7	49.0	48.9 \pm 6.3	50.0	0.716
Protein (%)	15.8 \pm 3.8	15.0	16.2 \pm 3.7	15.0	0.506
Fat (%)	34.8 \pm 6.8	34.0	34.8 \pm 5.7	33.0	0.884
SFA (g)	26.3 \pm 9.1	24.9	29.0 \pm 8.9	27.5	0.040
SFA (%)	12.7 \pm 4.0	12.02	13.6 \pm 3.7	12.69	0.027
Fiber (g)	20.6 \pm 7.1	20.0	18.5 \pm 6.5	18.5	0.083
Vitamin A (μg)	909.9 \pm 889.4	691.4	829.4 \pm 953.0	631.7	0.533
Vitamin E (mg)	13.3 \pm 5.7	12.1	12.5 \pm 5.2	11.9	0.566
Vitamin C (mg)	106.8 \pm 80.6	81.7	92.5 \pm 82.0	56.2	0.135
Thiamine (mg)	0.8 \pm 0.2	0.8	0.7 \pm 0.1	0.8	0.271
Riboflavin (mg)	1.4 \pm 0.5	1.4	1.6 \pm 0.4	1.5	0.065
Niacin (mg)	10.2 \pm 4.6	9.6	9.5 \pm 3.8	8.3	0.309
Vitamin B6 (mg)	1.2 \pm 0.3	1.2	1.1 \pm 0.3	1.1	0.707
Folate (μg)	114.9 \pm 33.6	114.1	114.7 \pm 36.7	116.7	0.964
Vitamin B12 (μg)	4.4 \pm 6.5	3.6	4.42 \pm 2.1	4.6	0.147
Sodium (mg)	4057.1 \pm 1381.	3918.9	4380.9 \pm 1384.7	4123.8	0.102
Potassium (mg)	2323.6 \pm 630.8	2271.1	2282.6 \pm 539.9	2221.8	0.771
Magnesium (mg)	252.5 \pm 63.5	241.3	245.9 \pm 47.3	240.3	0.754
Zinc (mg)	9.5 \pm 3.0	9.2	9.7 \pm 2.2	9.9	0.221
Iron (mg)	10.7 \pm 2.8	10.5	10.1 \pm 1.9	9.8	0.347

SFA: Saturated fatty acid; **PUFA:** Polyunsaturated fatty acid; **MUFA:** Monounsaturated fatty acid.
Mann-Whitney U test and Student's T test were used.

has been found to increase the rate of preferring high energy foods⁴². A negative relationship was found between amount of sleep and unhealthy eating habits⁴³. According to previous studies, the intake of fruits, vegetables and milk has a positive association, and the intake of sweets, snacks and fast food has negative association with sleep duration⁴⁴⁻⁴⁶. On the other hand, studies suggest that short sleep duration was related to increased fat intake^{47,48}. Poor sleep quality was associated with a lower intake of fruits, vegetables and milk and higher intake of soda, soft drinks, fast food, instant noodle and confectionaries⁴⁹. Similar to our result, in a study conducted on Iranian adolescents, there was no relationship between sleep quality and consumption of micronutrients and macronutrients. The mean intake of omega-3 fatty acids in subjects with good-

quality sleep was higher than that in the subjects with low-quality sleep⁵⁰.

In our study, it was observed that individuals who did not have regular breakfast had poorer sleep quality. However, not many studies have examined the association of sleep duration with skipping breakfast among adolescents⁵¹⁻⁵³. Among a large sample of Greek children and adolescents, insufficient sleep duration was associated with unhealthy dietary habits including skipping breakfast⁵¹. One study showed that skipping breakfast were significantly higher in children who reported poor sleep⁵². Skipping breakfast was associated with total and abdominal obesity in adolescents independent of sleep duration⁵⁴.

Prevalence of obesity has reached epidemic proportions across all gender, age and ethnic groups⁵⁵. Research to date in young children and adults shows a strong, inverse relationship between sleep duration and risk for overweight and obesity. Fewer studies examining this relationship have been conducted in adolescents⁵⁶. In this study, the prevalence of overweight and obesity was found to be 10.4% and 8.4%, respectively. Also, although statistical significance was not found, obese adolescents were shown to have less sleep duration averages than adolescents with normal body weight. In current study, BAZ of adolescents was also evaluated according to their sleep quality and no significant difference was found. It is thought that this result was obtained because more than half of the adolescents had normal body weight. In addition, a negative correlation was determined between the waist/hip ratio of students and the PSQI score. Sleep duration and quality have recently been described among obesity risk factors. Many epidemiological studies have shown that there is a relationship between less than 6-7 hours of sleep and obesity⁵⁷. However, causation is difficult to determine because of the fact that most of the epidemiological studies are observational⁵⁸. HELENA study results showed that European adolescents who slept 6 hours or less had a higher average BMI of 1.7kg/m² and 3.4cm larger waist circumference than those who slept 10 hours or more⁵⁹. In a study on adolescents who less than 8 hours had significantly higher body fat, waist and hip circumference, and BMI⁶⁰. Similar to our results, the findings of the Babu *et al.*'s study showed a nonmonotonic relationship between sleep quality and anthropometric parameters⁶¹.

There are some strengths and limitations of this study. To date, very few studies have investigated the relationship between dietary habits, food consumption, screen times, anthropometric measurements, and sleep quality in adolescents. We considered sleep duration and sleep quality in explaining sleep status. We considered numerous variables to investigate the association.

This was designed as a pilot study and therefore the sample chosen is not representative and the conclusions cannot be generalized. A single 24-hour recall is not considered to be representative of habitual diet. Further investigations should consider the relationship between sleep and related factors by using objective sleep measures such as polysomnography.

CONCLUSIONS

It was determined that most of the adolescents had good sleep quality and sleep duration is consistent with the recommended need. Sleep quality is affected by many factors such as dietary habits, food consumption, screen times, and individual factors. This study results support the development of interventions to help adolescents improve sleep quality. Further studies are needed to clarify the multiple mechanisms involved between individual factors, dietary habits, food consumption, anthropometric measurements, screen times, and sleep patterns in adolescents.

ACKNOWLEDGMENTS

We would like to thank the adolescents and their parents who devoted their time to the data collection.

AUTHORS' CONTRIBUTIONS

ÖK and GK designed the study, ARÖ collected the data, GK, ÖK, and ARÖ prepared the paper.

FUNDING

The authors have no financial relationships relevant to this article to disclose.

COMPETING INTERESTS

Authors state that there are no conflicts of interest in preparing the manuscript.

REFERENCES

- (1) Benedict C, Byberg L, Cedernaes J, Hogenkamp PS, Giedratz V, Kilander L, et al. Self-reported sleep disturbance is associated with Alzheimer's disease risk in men. *Alzheimers Dement*. 2015; 11(9): 1090-7.
- (2) Kronholm E, Laatikainen T, Peltonen M, Sippola R, Partonen T. Self-reported sleep duration, all-cause mortality, cardiovascular mortality and morbidity in Finland. *Sleep Med*. 2011; 12(3): 215-21.
- (3) Ramos AR, Jin Z, Rundek T, Russo C, Homma S, Elkind MS, et al. Relation between long sleep and left ventricular mass (from a multiethnic elderly cohort). *Am J Cardiol*. 2013; 112(4): 599-603.
- (4) Bayon V, Leger D, Gomez-Merino D, Vecchierini MF, Chennaoui M. Sleep dept and obesity. *Ann Med*. 2014; 46(5): 264-72.
- (5) Schmid SM, Hallschmid M, Schultes B. The metabolic burden of sleep loss. *Lancet Diabetes Endocrinol*. 2015; 3(1): 52-62.
- (6) Grandner MA, Chakravorty S, Perlis ML, Oliver L, Gurubhagavatula I. Habitual sleep duration associated with self-report and objectively determined cardiometabolic risk factors. *Sleep Med*. 2014; 15(1): 42-50.
- (7) Guo X, Zheng L, Wang J, Zhang X, Li J, Sun Y. Epidemiological evidence for the link between sleep duration and high blood pressure: a systematic review and meta-analysis. *Sleep Med*. 2013; 14(4): 324-32.
- (8) Paruthi S, Brooks LJ, D'Ambrosio C, Hall WA, Kotagal S, Lloyd RM, et al. Malow BA, Consensus statement of the American Academy of Sleep Medicine on the recommended amount of sleep for healthy children: methodology and discussion. *J Clin Sleep Med*. 2016; 12(11): 1549-61.
- (9) National Sleep Foundation. *Teens and Sleep*. (Access date 24 April 2020). Available at: <https://www.sleepfoundation.org/articles/teens-and-sleep>
- (10) Leger D, Bayon V, Sanctis A. The role of sleep in the regulation of body weight. *Mol Cell Endocrinol*. 2015; 418: 101-7.
- (11) Lytle LA, Pasch KE, Farbaksh K. The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring)*. 2011; 19(2): 324-31.
- (12) Miller MA, Kruisbrink M, Wallece J, Ji C, Cappuccio FP. Sleep duration and incidence of obesity in infants, children, and adolescents: a systematic review and meta-analysis of prospective studies. *Sleep*. 2018; 41(4): 1-19.
- (13) Koyuncuoğlu Güngör N. Overweight and Obesity in Children and Adolescents. *J Clin Res Pediatr Endocrinol*. 2014; 6(3): 129-43.
- (14) Chaput JP, Dutil C. Lack of sleep as a contributor to obesity in adolescents: impacts on eating and activity behaviors. *Int J Behav Nutr Phys Act*. 2016; 13(1): 103.
- (15) Kim S, DeRoo LA, Sandler DP. Eating patterns and nutritional characteristics associated with sleep duration. *Public Health Nutr*. 2011; 14(5): 889-95.
- (16) Buysse DJ, Reynolds CF 3rd, Monk TH, Berman SR, Kupfer DJ. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research. *Psychiatry Res*. 1989; 28(2): 193-213.
- (17) Agargün YM, Kara H, Anlar Ö. Pittsburg Uyku Kalitesi İndeksi'nin geçerliliği ve güvenilirliği. *Türk Psikiyatri Dergisi* 1996; 7(2): 107-15.
- (18) Gordon CC, Chumlea WC, Roche AF. Measurement descriptions and techniques Anthropometric Standardization Reference Manual. In: Lohman TG, Roche AF, Martorell R, eds. 1st edition, 3-12. Illinois: Kinetics Books Champaign; 1998.
- (19) de Onis M, Onyango AW, Borghi E, Siyam A, Nishida C, Siekmann J. Development of a WHO growth reference for school-aged children and adolescents. *Bull World Health Organ*. 2007; 85: 660-67.
- (20) World Health Organization. WHO Growth reference 5-19 years - application tools. (Access date 24 April 2020). Available at: <http://www.who.int/growthref/tools/en/>
- (21) Merdol Kutluay T. Toplu beslenme yapılan kurumlar için standart yemek tarifeleri. Ankara: Hatiboglu Yayınevi, Ankara, Turkey, 5th edition, 2015 (in Turkish).
- (22) Rakıcıoğlu N, Tek Acar, N, Ayaz A, Pekcan G. Photograph catalog of food and dishes: Portion sizes and amounts. Ata Ofset Pub, Ankara, Turkey, 2nd edition, 2009.
- (23) Ebspro for Windows, Stuttgart, Germany, BEBİS, Turkish version and Data Bases: Bundeslebensmittelschlüssel, 11.3 and other sources. Nutrition Information Systems, 2004.
- (24) IBM Corp. Released 2016. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.
- (25) World Health Organization (WHO). Sleep characteristics and sleep deprivation in infants, children and adolescents. WHO Regional Office for Europe, European Centre for Environment and Health Bonn Office: World Health Organization; 2004.
- (26) Xu Z, Su H, Zou Y, Chen J, Wu J, Chang W. Sleep quality of Chinese adolescents: distribution and its associated factors. *J Paediatr Child Health*. 2012; 48(2): 138-45.
- (27) Şenol V, Soyuer F, Pekşen Akça R, Argün M. The Sleep Quality in Adolescents and the Factors that Affect It. *Kocatepe Med J*. 2012; 14: 93-102.
- (28) Şimşek Y, Tekgöl N. Sleep quality in adolescents in relation to age and sleep-related habitual and environmental factors. *J Pediatr Res*. 2019; 6(4): 307-13.
- (29) Gradisar M, Gardner G, Dohnt H. Recent worldwide sleep patterns and problems during adolescence: A review and meta-analysis of age, region, and sleep. *Sleep Med*. 2011; 12: 110-8.
- (30) Felden ÉP, Filipin D, Barbosa DG, Andrade RD, Meyer C, Louzada FM. Fatores associados à baixa duração do sono em adolescentes. *Rev Paul Pediatr*. 2015; 34: 64-70.
- (31) Iglowstein I, Jenni OG, Molinari L, Largo RH. Sleep duration from infancy to adolescence: reference values and generational trends. *Pediatrics*. 2003; 111: 302-7.
- (32) Carskadon MA, Wolfson AR, Acebo C, Tzischinsky O, Seifer R. Adolescent sleep patterns, circadian timing, and sleepiness at a transition to early school days. *Sleep*. 1998; 21: 871-81.
- (33) Giannotti F, Cortesi F, Sebastiani T, Ottaviano S. Circadian preference, sleep, and daytime behavior in adolescence. *J Sleep Res*. 2002; 11: 191-99.
- (34) Tonetti L, Fabbri M, Natale V. Sex differences in sleep-time preference, and sleep need: a cross-sectional survey among Italian pre-adolescents, adolescents, and adults. *Chronobiol Int*. 2008; 25: 745-59.
- (35) Akçay D, Akçay BD. The influence of media on the sleep quality in adolescents. *Turk J Pediatr*. 2018; 60: 255-63.

- (36) Mei X, Zhou Q, Li X, Jing P, Wang X, Hu Z. Sleep problems in excessive technology use among adolescent: a systemic review and meta-analysis. *SSP*. 2018; 2: 9.
- (37) Garmy P, Nyberg P, Jakobsson U. Sleep and television and computer habits of Swedish school-age children. *J Sch Nurs*. 2012; 28: 469-76.
- (38) Van den Bulck J. Television viewing, computer game playing, and Internet use and self-reported time to bed and time out of bed in secondary-school children. *Sleep*. 2004; 27: 101-04.
- (39) American Academy of Pediatrics Committee on Public Education. Policy statement-Media education. *Pediatrics*. 2010; 126: 423-26.
- (40) Weiss A, Xu F, Storfer-Isser A, Thomas A, Ievers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep*. 2010; 33: 1201-9.
- (41) Grandner MA, Kripke DF, Naidoo N, Langer RD. Relationships among dietary nutrients and subjective sleep, objective sleep, and napping in women. *Sleep Med*. 2010; 11: 180-4.
- (42) Chaput JP. Sleep patterns, diet quality and energy balance. *Physiol Behav*. 2014; 134: 86-91.
- (43) Chaput JP, Katzmarzyk PT, LeBlanc AG, Tremblay MS, Barreira TV, Broyles ST, et al. Associations between sleep patterns and lifestyle behaviors in children: an international comparison. *Int J Obes Suppl*. 2015; 5(Suppl 2): S59-S65.
- (44) Ferranti R, Marventano S, Castellano S, Giogianni G, Nolfo F, Rametta S, et al. Sleep quality and duration is related with diet and obesity in young adolescent living in Sicily, Southern Italy. *Sleep Sci*. 2016; 9(2): 117-22.
- (45) Gong QH, Li H, Zhang XH, Zhang T, Cui J, Xu GZ. Associations between sleep duration and physical activity and dietary behaviors in Chinese adolescents: results from the Youth Behavioral Risk Factor Surveys of 2015. *Sleep Med*. 2017; 37: 168-73.
- (46) Kruger AK, Reither EN, Peppard PE, Krueger PM, Hale L. Do sleep-deprived adolescents make less-healthy food choices? *Br J Nutr*. 2014; 111(10): 1898-904.
- (47) Fisher A, McDonald L, van Jaarsveld CHM, Llewellyn C, Fildes A, Schrempft S, et al. Sleep and energy intake in early childhood. *Int J Obes*. 2014; 38(7): 926-29.
- (48) Weiss A, Xu F, Storfer-Isser A, Thomas A, Ievers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep*. 2010; 33(9): 1201-9.
- (49) Min C, Kim HJ, Park IS, Park B, Kim JH, Sim S, et al. The association between sleep duration, sleep quality, and food consumption in adolescents: A cross-sectional study using the Korea Youth Risk Behavior Web-based Survey. *BMJ Open*. 2018; 8: e022848.
- (50) Javadi M, Alimoradi F, Avani A, Jalilolghadr S. Association between sleep quality and intake of macronutrients and micronutrients in adolescents. *J Mazandaran Univ Med Sci*. 2018; 27(156): 205-10.
- (51) Tambalis KD, Panagiotakos DB, Psarra G, Sidossis LS. Insufficient sleep duration is associated with dietary habits, screen time, and obesity in children. *J Clin Sleep Med*. 2018; 14: 1689-96.
- (52) Agostini A, Lushington K, Kohler M, Dorrian J. Associations between self-reported sleep measures and dietary behaviours in a large sample of Australian school students (n=28,010). *J Sleep Res*. 2018; 27(5): e12682.
- (53) Smith KJ, Breslin MC, McNaughton SA, Gall SL, Blizzard L, Venn AJ. Skipping breakfast among Australian children and adolescents; findings from the 2011-12 National Nutrition and Physical Activity Survey. *Aust NZ J Public Health*. 2017; 41: 572-78.
- (54) Forkert ECO, Moraes ACFD, Carvalho HB, Manios Y, Widhalm K, Gonzales-Gross M, et al. Skipping breakfast is associated with adiposity markers especially when sleep time is adequate in adolescents. *Sci Rep*. 2019; 9: 6380.
- (55) Di Renzo L, Tyndall E, Gualtieri P, Carboni C, Valente R, Ciani AS, et al. Association of body composition and eating behavior in the normal weight obese syndrome. *Eat Weight Disord*. 2016; 21(1): 99-106.
- (56) Lytle LA, Pasch KE, Farbaksh K. The relationship between sleep and weight in a sample of adolescents. *Obesity (Silver Spring)*. 2011; 19(2): 324-31.
- (57) Patel SR, Hu FB. Short sleep duration and weight gain: a systematic review. *Obesity (Silver Spring, Md)*. 2008; 16(3): 643-53.
- (58) S Bruce E, Lunt L, E McDonagh J. Sleep in adolescents and young adults. *Clin Med*. 2017; 17(5): 424-8.
- (59) Garaulet M, Ortega FB, Ruiz JR, Rey-Lopez JP, Beghin L, Manios Y, et al. Short sleep duration is associated with increased obesity markers in European adolescents: effect of physical activity and dietary habits. The HELENA study. *Int J Obes*. 2011; 35(10): 1308-17.
- (60) Weiss A, Xu F, Storfer-Isser A, Thomas A, Ievers-Landis CE, Redline S. The association of sleep duration with adolescents' fat and carbohydrate consumption. *Sleep*. 2010; 33(9): 1201-9.
- (61) Babu R, Bahuleyan B. Correlation of sleep quality with anthropometric parameters in young healthy individuals. *Int J Res Med Sci*. 2018; 6: 613-7.