

# Revista Española de Nutrición Humana y Dietética

## Spanish Journal of Human Nutrition and Dietetics



CrossMark  
click for updates

www.renhyd.org



### RESEARCH ARTICLE

## Assessment of the glycemic index in a white bread enriched with fermented cassava dough using *Lactobacillus plantarum* in apparently healthy volunteers

R Marisol Martínez<sup>a</sup>, Mario Cruz<sup>b</sup>, Araceli Loredó-Treviño<sup>a</sup>, José Luis Martínez-Hernández<sup>a</sup>,  
Aurora Garza<sup>c</sup>, Ruth Belmares<sup>a,\*</sup>

<sup>a</sup> Departamento de Investigación en Alimentos, Facultad de Ciencias Químicas, Universidad Autónoma de Coahuila, Saltillo, Coahuila, México.

<sup>b</sup> Departamento Ciencia y Tecnología en Alimentos, Universidad Autónoma Agraria Antonio Narro, Buena Vista, Coahuila, México.

<sup>c</sup> Departamento de Bioquímica y Medicina Molecular, Facultad de Medicina, Universidad Autónoma de Nuevo León, Nuevo León, México.

\*[ruthbelmares@uadec.edu.mx](mailto:ruthbelmares@uadec.edu.mx)

Assigned Editor: Édgar Pérez Esteve. Universitat Politècnica de València, Valencia, Spain.

Received: 10/18/2023; Accepted: 03/07/2024; Published: 05/21/2024.

#### KEYWORDS

Glycemic Index,  
Cassava flour;  
Bread;  
Fermentation.

#### ➤ Assessment of the glycemic index in a white bread enriched with fermented cassava dough using *Lactobacillus plantarum* in apparently healthy volunteers

#### ABSTRACT

**Introduction:** The substitution of wheat flour with cassava flour has resulted in high glycemic indices or products with low quality, while limited research on substitution with fermented dough has shown improved quality. Therefore, the aim of this research was to evaluate the In vivo glycemic index of a baked product supplemented with cassava dough fermented with lactic acid bacteria.

**Methodology:** Twenty-eight young adult students both genders participated, of whom 15 met the stipulated standards. Overall and by gender, the glycemic index of bread enriched to 20% with fermented cassava dough with glucose as standard was determined by capillary glycemia.

**Results:** There was a significant difference in the glycemic curve of bread against the standard. An average glycemic index of  $40 \pm 6.33$  was obtained, classifying it as "Low", showing a different index between genders of  $52 \pm 8.21$  in men and  $36 \pm 6.96$  in women with no significant difference.

**Conclusions:** The use of fermented cassava dough as a partial substitution in wheat-based baked products is favorable for obtaining products with a lower glycemic index.

**Funding:** This research was supported by the CONACYT scholarship program in collaboration with the Universidad Autónoma de Coahuila, Faculty of Chemical Sciences, Departamento de Investigación en Alimentos, Functional Foods & Nutrition, with scholarship number 1082548. Additionally, support was provided by the Department of Food Science and Technology, Universidad Autónoma Agraria Antonio Narro.



## PALABRAS CLAVE

Índice Glicémico;  
Harina de Yuca,  
Pan;  
Fermentación.

➤ **Determinación del índice glucémico en pan blanco enriquecido con masa de yuca fermentada con *Lactobacillus plantarum* en voluntarios aparentemente saludables**

**RESUMEN**

**Introducción:** La sustitución de harina de trigo por harina de yuca ha resultado en índices glicémicos altos o productos con baja calidad, mientras que las limitadas investigaciones de sustitución con masa fermentada muestran una mejor calidad, por lo que el objetivo de esta investigación fue evaluar *in vivo* el índice glicémico de un producto horneado adicionado con masa de yuca fermentada con bacterias ácido-lácticas.

**Metodología:** Participaron 28 estudiantes adultos jóvenes de ambos sexos de los cuales se aceptaron 15 bajo los estándares estipulados. En conjunto y por género, se determinó por glicemia capilar el índice glicémico de un pan enriquecido al 20% con masa fermentada de yuca con glucosa como estándar.

**Resultados:** Se presentó diferencia significativa en la curva glicémica del pan contra el estándar. Se obtuvo  $40 \pm 6,33$  de índice glicémico promedio en el pan clasificándolo como "Bajo", mostrando un índice muy diferente entre el género de  $52 \pm 8,21$  en hombres y  $36 \pm 6,96$  en mujeres sin diferencia significativa.

**Conclusiones:** El empleo de masa fermentada de harina de yuca como sustitución parcial en producto horneado a base de trigo es favorable en la obtención de productos con menor índice glicémico.

**Financiación:** Esta investigación fue financiada como parte del programa de becas CONACYT en conjunto con la Universidad Autónoma de Coahuila Facultad de Ciencias Químicas, Departamento de Investigación de Alimentos, *Functional Foods & Nutrition*, con número de beca 1082548. Así como el Departamento de Ciencia y Tecnología en Alimentos, Universidad Autónoma Agraria Antonio Narro.

KEY  
MESSAGES

1. White bread with a 20% substitution of cassava flour fermented dough has a higher protein and lipid content, as well as a lower amount of carbohydrates in its nutritional composition than commercial bread.
2. The glycemic index of the white bread in the study on average is classified as low.
3. Although the classification of the index is maintained when analyzed by gender, variations occur, showing higher values in men than in women.
3. The use of sourdough based on cassava flour as a substitute is favorable for achieving a low glycemic index.

## CITATION

Martínez RM, Cruz M, Loredó-Treviño A, Martínez-Hernández JL, Garza A, Belmares R. Assessment of the glycemic index in a white bread enriched with fermented cassava dough using *Lactobacillus plantarum* in apparently healthy volunteers. Rev Esp Nutr Hum Diet. 2024; 28(2): 93-101.  
doi: <https://doi.org/10.14306/renhyd.28.2.2053>

## INTRODUCTION

The glycemic index (GI) is defined as the comparison of the area under the glycemic response curve following the ingestion of a single food. The higher this index, the greater the ease and speed with which glucose from the food is digested and absorbed. Literature reveals that the consumption of low-GI foods provides health benefits; therefore, researchers seek to modify the GI of staple foods around the world, with bread being one of the most significant staple foods<sup>1,2</sup>.

Bread is the most consumed cereal-based food worldwide, typically made from wheat flour (WF) with a GI ranging from moderate to high. However, the approach has emerged to use non-traditional flour sources that are resilient to changing climate conditions, are more cost-effective, and contribute to enhancing food security for the population. One such example is the use of cassava tubers<sup>3,4</sup>.

Cassava or Manihot, is a high-yielding drought-resistant crop considered a staple food as it serves as a subsistence crop. Its use in the form of cassava flour (CF) has garnered interest due to its excellent stability, extraordinary thickening capacity, and lower retrogradation and gelatinization indices. Previously, the use of CF has been described as a partial substitute for baked products based on WF, Okafor *et al.* in 2018<sup>5</sup>, and its evaluation of GI in bar-type breads with different levels of substitution resulted in a GI range of 91-94, which is higher than the average. However, a viable approach related to reducing the GI and improving the properties of baked products is the fermentation of the dough<sup>6-8</sup>.

Fermentation of dough provides several benefits, including improvement in the aroma, flavor, texture, shelf life, an increase in short-chain essential fatty acids (SCFA) and a reduction in GI in the baked product<sup>9,10</sup>. The use of non-traditional flour sources as a base in obtaining fermented dough as a partial substitute for recipes using WF can be encouraging for the innovation of staple products, making use of accessible and cost-effective crops. Therefore, the objective of this research was to evaluate the GI of a white bread with partial substitution of 20% CF sourdough to classify its impact on apparently healthy volunteers.

## METHODOLOGY

This project underwent review by the institutional scientific ethics committee for scientific development at the Faculty of Chemical Sciences, *Universidad Autónoma de Coahuila*, and was approved under registration number TDCYTA-20-10-22-1. All volunteers

were informed of the protocol orally and through a printed informed consent in compliance with the Sanitary Control of Products and Services Regulation<sup>11</sup>, the General Health Law Regulation on Health Research<sup>12</sup>, as well as ISO 26642:2012<sup>13</sup> pertaining to the protocol for determining the GI. This research employed a quantitative, non-experimental, and cross-sectional descriptive design.

### Baked product

The food under study was a white bread with a 20% substitution of cassava fermented dough (CFD) fermented for 32 hours at 36 °C with *Lactobacillus plantarum*, which was obtained from the reservoir of microorganisms belonging to the Research Department of Food. The bread was obtained under a standard recipe (WF: 320 g, CFD: 80 g, milk: 250 mL, sugar: 15 g, salt: 7 g, unsalted butter: 50 g, instant dry yeast: 12 g) with three kneading cycles, including a 30-minutes rest period between kneading cycles to stimulate autolysis. Subsequently, a nutritional analysis was conducted to determine its bromatological profile: Moisture, ash content, fat content (Soxhlet), protein content (Kjeldahl/conversion factor 6.25), crude fiber content was determined according to the AOAC methods<sup>14</sup>; total carbohydrate content was determined by difference and Total Sugar Content by Phenol-Sulfuric Acid method described by DuBois<sup>15</sup>.


### Volunteers

Due to the variability of GI influenced by several factors, of a total of 25 apparently healthy volunteers from the *Universidad Autónoma de Coahuila*, aged between 20 and 40 years, who participated in this study, only 18 volunteers were selected based on the acquisition and analysis of ABCD data shown in Figure 1. A SD Lipid Care® Analyzer (Brand: SD Biosensor, INC, Elaborado: Kana Undesa S.A. de C.V., México) was used to measure HDL cholesterol, LDL cholesterol, total cholesterol, total triglycerides, and capillary blood glucose levels. Following this, a brief questionnaire about lifestyle changes and physical activity was applied and a 2000 kcal dietary plan was implemented for five days to standardize carbohydrate consumption before the test. On three separate occasions, volunteers were requested to fast for 6-8 hours before each analysis.

### GI evaluation

Following the protocol specified by FAO/WHO and ISO<sup>13,16</sup>, capillary blood samples were taken to obtain glucose levels at 0, 15, 30, 45, 60, 90, and 120 minutes after the consumption of the standard and white bread-cassava. Due to the variations that commercial white bread may have in its nutritional content (raw material and processes), a valued and stable oral glucose tolerance solution was selected to determine glucose

**Figure 1.** Data ABCD collected for volunteer selection.



tolerance (Brand: Glucox, Manufactured: Materiales y Abastos Especializados S.A. de C.V., Mexico). The GI was obtained using Equation [1]:

$$GI\% = \frac{IAUC\ bread}{IAUC\ glucose} * 100 \quad \text{Eq [1]}$$

Where:

IAUC= Incremental area under the blood glucose response curve.

### Statistical analysis

The data obtained from the accepted volunteers for this study was divided into females (F), males (M), and total data (T) to conduct a more in-depth analysis of the collected data. All results are reported as mean values with standard deviation. To determine significant differences ( $p < 0.05$ ), a means comparison was performed using the Tukey test at 95% confidence interval.

## RESULTS

Enough white bread was made to use the same batch in all analyses. The nutritional composition of white bread with fermented cassava dough (mentioned as white bread-cassava in the following) are presented in Table 1, where a very different profile is observed from its commercial version (Protein: 8.2 g/100 g, Lipid: 3 g/100 g and Carbohydrates: 50.4 g/100 g<sup>17</sup>). Continuing with the analysis of glycemic index, the average age in this study was 27 years, with a predominance of female participants.

**Table 1.** Nutritional composition (mean±standard deviation) of white bread-cassava.

	g/100 g
Moisture	31.78±0.97
Ash	5.22±0.71
Lipids	10.38±0.02
Protein	15.17±0.5
Fiber	0.11±0.007
Total carbohydrates	37.34±0.86
Total sugars	9±0.85

Anthropometric results, in general, indicate a normal BMI (23.96±3.26 kg/m<sup>2</sup>). However, M (25.22±3.51 kg/m<sup>2</sup>) showed slightly abnormal values compared to F (23.61±3.23 kg/m<sup>2</sup>), primarily due to the average muscle percentage (F: 25.57±2.74%, M: 33.03±5.49%, T: 27.23±4.61%). Likewise, the waist-to-hip ratio (WHR) values fall within the normal range (0.78±0.06). On the other hand, the biochemical results are presented in Table 2, with values within the established standards. These data provide an overview of the volunteers' health status, classifying them as healthy and fit for this study.

Clinical and Dietary data reveal interesting insights. In the Personal medical history aspect, it is observed that 67% of the volunteers had experienced COVID-19, and 83% of them changed their habits due to the pandemic. In the Familiar medical history, 72% of the volunteers have family members with type 2 diabetes, 67% hypertension, 39% some type of cancer and 22% hypothyroidism. On the Dietary data, meal timings are quite similar (T: 3±1), and water intake is deficient (T: 1.8 L). In terms of sleep duration, F (8±1.4) sleep 2 hours more on average than M (6±0.5), categorizing them differently in terms of sleep patterns<sup>18</sup>. The Food frequency reported by the volunteers shows high consumption of both recommended and non-recommended foods according to the ENSANUT CONTINUA 2022 classification. The ABCD data collected and displayed encompass 18 volunteers who passed the biochemical phase and, therefore, had their data included in this part of the study's statistics.

Primarily, Figure 2 shows the increase in glycemia caused by glucose and white bread-cassava. This phase includes the final group of 15 volunteers who met the requirements on the days of the GI analysis. Each volunteer received a portion of 133.9±0.4 g of bread for the curve. We observed a lower increase in glucose with bread, with a much more prolonged and sustained decline, staying below reference values with glucose.

On the other hand, glucose exhibited an intermediate peak that gradually decreased, showing a significant difference from 30 to 60 minutes and again at 120 minutes. On average, a GI of 40 was obtained, classifying it as low, as shown in Table 3.

## DISCUSSION

The volunteers present an adequate nutritional status, with a high prevalence of diseases in their family medical history and significant changes in their habits, including decreased physical activity and irregular sleep patterns. According to food frequency, the duality presented by the young individuals in their efforts to maintain a healthy lifestyle through good


**Table 2.** Biochemical data (mean±standard deviation) related to the previous health status of the volunteers.

	mg/dL	Female	Male	Total
<b>Biochemical data</b>	Triglycerides	90.7±51.6	127.8±39.3	98.9±50.6
	Total cholesterol	147±31.3	144.8±34.8	146.5±31.1
	LDL cholesterol	86.3±29	79±39.2	84.6±30.4
	HDL cholesterol	42.3±8.4	40.3±8.2	41.8±8.2
	Glucose	88.4±6.6	93±6.4	89.4±6.7

nutrition and the ease of obtaining ready-to-eat products that provide convenience and comfort are part of the new normal. Overall, the white bread-cassava is in the Low GI classification and this classification remains consistent when separating them by gender, despite variations that may be attributable to various factors, both modifiable and non-modifiable.

Fang *et al.*, 2023<sup>19</sup>, mentions that flour fermentation in general, reduces the GI of bread derived from the production of organic acids and from the important reduction in the digestibility of native starch, also explains that wheat breads with sourdough have low GI, and also mentions that sourdough type II (dough fermented with a starter culture) has advantages

**Figure 2.** Glycemic response following the ingestion of oral glucose and white bread-cassava.



**Table 3.** GI of white bread-cassava.

	n	GI (Glucose=100 g)	Classification <sup>1</sup>
<b>Male</b>	4	52±8.21 <sup>a</sup>	Low
<b>Female</b>	11	36±6.96 <sup>a</sup>	Low
<b>Total</b>	15	40±6.33 <sup>a</sup>	Low

<sup>1</sup>GI High: ≥70; GI Medium: 56 a 69; IG Low: ≤55.

Data are expressed as mean±standard deviation.

<sup>2</sup>Values with different letters within the same column are significantly different (p<0.05).

such as better control of fermentation parameters and better nutritional properties. Garrido-Galand *et al.*, 2021<sup>20</sup>, concluded that fermenting legume flour enhances techno-functional properties, while fermenting cereal flour increase antioxidant properties. Dough fermentation endows desirable nutritional and rheological properties to baked products, notable properties in cassava-white bread.

Oyeyinka *et al.*, 2020<sup>21</sup>, mentioned that fermentation in cassava is used to improve its functionality, being acid-lactic bacteria responsible for these processes. Longoria *et al.*, 2020<sup>3</sup>, conducted a study using cassava dough fermented with *L. casei*, concluding that fermentation in short periods improves the rheological properties of non-traditional flour-based products. Penido *et al.*, 2018<sup>22</sup>, concluded that in spontaneous fermentations with bitter cassava, acid-lactic bacteria prevail throughout the process, being *L. plantarum* one of the species present and with better performance as starter culture. For this reason, selecting *L. plantarum*, a positive change in the composition of the bread was expected and this is demonstrated in its nutritional profile, very different from traditional white bread.

Is evident that the quality and quantity of carbohydrates in the food are crucial in determining the GI. Additionally, the interactions that starch has with other components affect its availability. Luhovyy *et al.*, 2022<sup>23</sup>, explain that 7 to 10 grams of protein are required to observe a hypoglycemic effect, of course it depends on the type of protein and the overall composition of the food. Additionally, starch can interact with lipids, phytochemicals, proteins, and hydrocolloids, and these mechanisms can interfere with starch metabolism, reducing its availability<sup>24-26</sup>.

The bread in this project exhibited good rheological properties and a nutritional profile different from the traditional one but with a favorable GI. The bread in this project exhibited good

rheological properties and a nutritional profile different from the traditional type but with a favorable GI. These attributes can be attributed to the recipe, the bread-making process, and the addition of fermented cassava dough, which is mainly responsible for the significant increase in protein content and the reduction of total carbohydrates. Okafor *et al.*, 2017<sup>5</sup>, stipulated that the addition of CF to the standard WF white bread recipe produces a high GI, however, in this study it was demonstrated that a previous fermentation with one of the naturally predominant BAL of CF, modifies its nutritional profile due to the decrease of free sugar content, increase of organic acids and reduction of starch digestibility<sup>19</sup>.

The white bread-cassava, both on average and by gender, is in the classification of low glycemic index, however, although there is no significant difference between them, the difference between the genders is remarkable. After the beginning of the strategy for an adequate reopening of activities of social, educational and economic nature called "The new normality"; the changes and new norms that it brought with it modified the social psyche because according to the National Epidemiologic Risk Symphor, the activities were or were not allowed<sup>27</sup>. In general, the factors with which the individual interacts have repercussions on the habits and therefore on the state of health in general, which includes metabolism.

Jarvis *et al.*, 2023<sup>28</sup>, decree that experiencing consecutive spikes in glucose levels leads to increased hunger sensations as well as lower mental health and sleep quality. Sleep is one of these factors and is crucial in regulating the endocrine and metabolic systems. Mohammadi *et al.*, 2021<sup>29</sup>, concluded that the duration of sleep does not directly interfere with the GI, however, they found a positive relationship between glycemic load and longer sleep duration. Ayer *et al.*, 2022<sup>30</sup>, conducted a study on adults in Brazil and found that more than half of the population modified the number of meals they consumed, of which 1/3 increased the number of meals. Additionally, the same pattern of increase in recommended and non-recommended food was presented, with a particular rise in the consumption of leafy green vegetables and bakery products. It is interesting to observe the current habits, as although they may be non-specific, they do show a certain relationship with the GI.

## CONCLUSIONS

The addition of cassava fermented dough to the standard formulation of a traditional wheat flour white bread decreases the average GI, making it favorable to use fermented dough as a partial or total replacement of a bread with a lower GI.

## ACKNOWLEDGMENTS

We thank Chemistry María García Braham for her leadership in sample collection and management of biological waste.

## AUTHORS' CONTRIBUTIONS

R.M.M. was the main researcher, an important part of sample collection and management of biological waste, interpretation of results and the main writer of this article. R.B. contributed to the creation and design of the study, supervised compliance with the established protocol and performed data analysis with the work team. M.C. contributed to the creation of the statistical design and interpretation of results, as well as providing financing from the *Universidad Autónoma Agraria Antonio Narro*. J.L.M. and A.L. contributed to the collection of data for the approval of the protocol before the ethics committee and performed data analysis with the work team. A.G. consulted with the ethics committee and supervised compliance with the established protocol.

All authors critically reviewed this and previous versions of the document.

## FUNDING

The *Consejo Nacional de Ciencia y Tecnología* (CONACYT) of Mexico provides financing by granting a postgraduate scholarship to the student Rosa Marisol Martínez Hernández, with scholarship number: 1082548. As well as internal resources from the *Universidad Autónoma de Coahuila*, Faculty of Chemical Sciences, Departamento de Investigación en Alimentos, Functional Foods & Nutrition. Additionally, support was provided by the Department of Food Science and Technology, *Universidad Autónoma Agraria Antonio Narro*.

## COMPETING INTERESTS

The authors express that there are no conflicts of interest when writing the manuscript.

## REFERENCES

- (1) Atkinson FS, Brand-Miller JC, Foster-Powell K, Buyken AE, Goletzke J. International tables of glycemic index and glycemic load values 2021: a systematic review. *Am J Clin Nutr.* 2021; 114(5): 1625-32. doi: 10.1093/ajcn/nqab233.
- (2) Stamatakis NS, Yanni AE, Karathanos VT. Bread making technology influences postprandial glucose response: A review of the clinical evidence. *Br J Nutr.* 2017: 1001-12. doi: 10.1017/S0007114517000770.
- (3) Longoria S, Contreras J, Belmares R, Cruz M, Flores M. Effect of Short Fermentation Times with *Lactobacillus paracasei* in Rheological, Physical and Chemical Composition Parameters in Cassava Dough and Biscuits. *Appl Sci.* 2020; 10(4): 1383. doi: 10.3390/app10041383.
- (4) Tiefenbacher KF Glossary of Terms in Wafers, Waffles and Adjuncts. *The Technology of Wafers and Waffles II.* Elsevier; 2019. p. 325-411. doi: 10.1016/B978-0-12-809437-2.00010-1
- (5) Okafor EN, Erukainure OL, Ozumba AU, Adewale CO, Kayode FO, Asieba GO, et al. Cassava Flour Substitution Modulates Glycemic Responses and Glycemic Index of Wheat Breads in Apparent Healthy Volunteers. *J Diet Suppl.* 2017; 14(4): 446-52. doi: 10.1080/19390211.2016.1267061.
- (6) Chisenga SM, Workneh TS, Bultosa G, Alimi BA, Siwela M. Dough rheology and loaf quality of wheat-cassava bread using different cassava varieties and wheat substitution levels. *Food Biosci.* 2020; 34: 100529. doi: 10.1016/j.fbio.2020.100529.
- (7) Cock JH, Connor DJ. Cassava. *Crop Physiology Case Histories for Major Crops.* Elsevier; 2021. p. 588-633. doi: 10.1016/B978-0-12-819194-1.00019-0.
- (8) Aryeetey E, Wireko-Manu FD, Asante JO, Laryea D, Ellis WO, Oduro I, et al. Recipe standardization of bread using cassava-wheat composite flour. *Journal of Culinary Science & Technology.* 2019; 17(3): 232-55. doi: 10.1080/15428052.2018.1429972.
- (9) Roobab U, Batool Z, Manzoor MF, Shabbir MA, Khan MR, Aadil RM. Sources, formulations, advanced delivery and health benefits of probiotics. *Curr Opin Food Sci.* 2020; 32: 17-28. doi: 10.1016/j.cofs.2020.01.003.
- (10) Papadimitriou K, Zoumpopoulou G, Georgalaki M, Alexandraki V, Kazou M, Anastasiou R, et al. Sourdough Bread. *Innovations in Traditional Foods.* Elsevier; 2019. p. 127-58.
- (11) Secretaría de Salud REGLAMENTO de Control Sanitario de Productos y Servicios. Obtenido de REGLAMENTO de Control Sanitario de Productos y Servicios. <http://www.conacyt.mx/cibiogem/images/cibiogem/normatividad/vigente/SALUD/071005RCSPS.pdf>.
- (12) Secretaría de Salud REGLAMENTO de la Ley General de Salud en Materia de Investigación para la Salud. Obtenido de REGLAMENTO de la Ley General de Salud en Materia de Investigación para la Salud: <http://www.conacyt.mx/cibiogem/images/cibiogem/normatividad/vigente/SALUD/071006RLSMIS.pdf>.
- (13) ISO 26642:2010(E) Food products – Determination of the glycaemic index (GI) and recommendation for food classification. ISO 2010. 2010:1-18.
- (14) AOAC International Official Method of Analysis. Post revised on date 15/03/2021. 1980.
- (15) DuBois Michel, Gilles KA, Hamilton JK, Rebers PA, Smith Fred



- Colorimetric Method for Determination of Sugars and Related Substances. Anal Chem. 1956; 28(3): 350-6. doi: 10.1021/ac60111a017.
- (16) FAO/WHO Report of a Joint FAO/WHO Expert Consultation. Carbohydrates in human nutrition., vol. 66. Rome: FAO Food and Nutrition Paper-66; 1997. p. 25-30.
- (17) Pérez A, Castro A, Palacios B, Flores I SMAE. 4.a edición. Mexico: ISBN 978-607-00-7928-3; 2014.
- (18) Andrea Contreras S. Sueño a lo largo de la vida y sus implicancias en salud. Revista Médica Clínica Las Condes. 2013; 24(3): 341-9. doi: 10.1016/S0716-8640(13)70171-8.
- (19) Fang L, Wang W, Dou Z, Chen J, Meng Y, Cai L, et al. Effects of mixed fermentation of different lactic acid bacteria and yeast on phytic acid degradation and flavor compounds in sourdough. LWT. 2023; 174: 114438. doi: 10.1016/j.lwt.2023.114438.
- (20) Garrido-Galand S, Asensio-Grau A, Calvo-Lerma J, Heredia A, Andrés A. The potential of fermentation on nutritional and technological improvement of cereal and legume flours: A review. Food Res Int. 2021; 145: 110398. doi: 10.1016/j.foodres.2021.110398.
- (21) Oyeyinka SA, Adeloye AA, Olaomo OO, Kayitesi E. Effect of fermentation time on physicochemical properties of starch extracted from cassava root. Food Biosci. 2020; 33: 100485. doi: 10.1016/j.fbio.2019.100485.
- (22) Penido FCL, Piló FB, Sandes SH de C, Nunes AC, Colen G, Oliveira E de S, et al. Selection of starter cultures for the production of sour cassava starch in a pilot-scale fermentation process. Braz J Microbiol. 2018; 49(4): 823-31. doi: 10.1016/J.BJM.2018.02.001.
- (23) Luhovyy BL, Kathirvel P. Food proteins in the regulation of blood glucose control. Adv Food Nutr Res. 2022; 102: 181-231. doi: 10.1016/bs.afnr.2022.05.001.
- (24) Wang S, Chao C, Cai J, Niu B, Copeland L, Wang S. Starch-lipid and starch-lipid-protein complexes: A comprehensive review. Compr Rev Food Sci Food Saf. 2020; 19(3): 1056-79. doi: 10.1111/1541-4337.12550.
- (25) Mahmood K, Kamilah H, Shang PL, Sulaiman S, Ariffin F, Alias AK A review: Interaction of starch/non-starch hydrocolloid blending and the recent food applications. Food Biosci. 2017; 19: 110-20. doi: 10.1016/j.fbio.2017.05.006.
- (26) Yang C, Zhong F, Douglas Goff H, Li Y Study on starch-protein interactions and their effects on physicochemical and digestible properties of the blends. Food Chem. 2019; 280: 51-8. doi: 10.1016/j.foodchem.2018.12.028.
- (27) Centeno J-E. LA NUEVA "NORMALIDAD" EN MÉXICO. Revista de Salud Publica. 2020; 87-90. doi: 10.31052/1853.1180.v0.n0.28963.
- (28) Jarvis PRE, Cardin JL, Nisevich-Bede PM, McCarter JP. Continuous glucose monitoring in a healthy population: understanding the post-prandial glycemic response in individuals without diabetes mellitus. Metabolism. 2023; 146: 155640. doi: 10.1016/j.metabol.2023.155640.
- (29) Mohammadi M, Nadjarzadeh A, Mirzaei M, Fallahzadeh H, Haghghatdoost F, Sakhaei R, et al. Dietary glycemic index and glycemic load in association with sleep duration: YaHS-TAMYZ and Shahedieh observational studies. Clin Nutr ESPEN. 2021; 46: 471-6. doi: 10.1016/j.clnesp.2021.09.007.
- (30) Ayer C, Celep AGS. Assessment of dietary habits and use of nutritional supplements in COVID-19: A cross-sectional study. PharmaNutrition. 2022; 22: 100309. doi: 10.1016/j.phanu.2022.100309.