

Revista Española de Nutrición Humana y Dietética

Spanish Journal of Human Nutrition and Dietetics



CrossMark
click for updates

www.renhyd.org



RESEARCH ARTICLE

Characterization of three varieties of Malaysian rambutan seed oil

Fatemeh Ghobakhlou^a, Hasanah Mohd Ghazali^{a,*}, Roselina Karim^a, Abdulkarim Sabo Mohammed^a

^aDepartment of Food Science, Faculty of Food Science and Technology, Universiti Putra Malaysia, Serdang, Malaysia.

*hasanah@upm.edu.my

Editor assigned: Miguel Ángel Lurueña. Comité Editorial de la Revista Española de Nutrición Humana y Dietética. Pamplona, España.

Received: 12/02/2019; accepted: 29/07/2019; published: 29/07/2019.

KEYWORDS

Rambutan;
Nephelium lappaceum;
Chemistry.

Characterization of three varieties of Malaysian rambutan seed oil

ABSTRACT

Introduction: Rambutan seed is considered a waste by-product of fruit processing. The seed waste is usually discarded or disposed in a large amount without an economic value which has become an issue that needs to be solved. However, the seed contains a considerable amount of crude fat. Aim: the purpose of this study was to characterize the physicochemical properties of oil of the seeds of three varieties of Malaysian rambutan fruit (R4, R7 and Serjan) for potential application.

Material and Methods: In this study, colour, refractive index, viscosity, free fatty acid content, peroxide value, p-anisidine value, iodine value, saponification value, unsaponifiable matter, fatty acid composition, thermal behaviour, melting point, and solid fat content of rambutan seed oil from three varieties were used to characterise physicochemical properties of rambutan seed oil.

Results: No significant differences ($p > 0.05$) were observed for free fatty acid content, peroxide value, p-anisidine value, saponification value, unsaponifiable matter, colour, viscosity and refractive index among three varieties. There are significant differences ($p < 0.05$) in the iodine value and melting point among the rambutan seed oil of three varieties. Varieties R7 and R4 seed had the highest ($37.62 \pm 0.10\%$) and lowest ($34.25 \pm 0.07\%$) crude fat content, respectively. Oleic ($37.75\text{--}40.58\%$) and arachidic ($35.24\text{--}36.89\%$) acids were the major fatty acids in the oil. Both melting and crystallization curves showed that the oil exhibited three distinct peaks. The complete melting and crystallization onset temperatures of the oil were $24.76\text{--}26.57^\circ\text{C}$ and $21.19\text{--}22.79^\circ\text{C}$, respectively.

Conclusions: This study reveals that the rambutan seed oil has potential to be used in various sectors of food industry. Therefore, rambutan seed can be fully utilized and, subsequently, the amount of waste can be minimized.

Caracterización de tres variedades de aceite de semillas de rambután de Malasia

PALABRAS CLAVE

Rambután;

Nephelium lappaceum;

Química.

RESUMEN

Introducción: La semilla de rambután se considera un subproducto residual del procesamiento del fruto. Los residuos de semillas generalmente se desechan o se eliminan en grandes cantidades sin un valor económico que se ha convertido en un problema que debe resolverse. Sin embargo, la semilla contiene una cantidad considerable de grasa cruda. Objetivo: el propósito de este estudio fue caracterizar las propiedades fisicoquímicas del aceite de las semillas de tres variedades de rambután de Malasia (R4, R7 y Serjan) para su posible aplicación.

Material y Métodos: En este estudio, el contenido de aceite de semilla de rambután de tres variedades se utilizó para caracterizar las propiedades fisicoquímicas del aceite de semilla de rambután en color, índice de refracción, viscosidad, contenido de ácidos grasos libres, índice de peróxido, índice de p-anisidina, índice de yodo, valor de saponificación, materia insaponificable, composición de ácidos grasos, comportamiento térmico, punto de fusión y grasa sólida.

Resultados: No se observaron diferencias significativas ($p > 0.05$) para el contenido de ácidos grasos libres, el valor de peróxido, el valor de p-anisidina, el valor de saponificación, la materia insaponificable, el color, la viscosidad y el índice de refracción entre tres variedades. Existen diferencias significativas ($p < 0.05$) en el valor de yodo y el punto de fusión entre el aceite de semilla de rambután de tres variedades. Las variedades R7 y R4 tuvieron el contenido de grasa cruda más alto ($37,62 \pm 0,10\%$) y más bajo ($34,25 \pm 0,07\%$), respectivamente. El ácido oleico ($37,75-40,58\%$) y el ácido araquídico ($35,24-36,89\%$) fueron los principales ácidos grasos en el aceite. Las curvas de fusión y cristalización mostraron que el aceite exhibía tres picos distintos. Las temperaturas de inicio de fusión y cristalización completas del aceite fueron $24,76-26,57^\circ\text{C}$ y $21,19-22,79^\circ\text{C}$, respectivamente.

Conclusiones: Este estudio revela que el aceite de semilla de rambután tiene potencial para ser utilizado en varios sectores de la industria alimentaria. Por lo tanto, la semilla de rambután se puede utilizar por completo y, en consecuencia, se puede minimizar la cantidad de desechos.

CITATION

Ghobakhlou F, Mohd Ghazali H, Karim R, Sabo Mohammed A. Characterization of three varieties of Malaysian rambutan seed oil. Rev Esp Nutr Hum Diet. 2019; 23(3): 183-93. doi: 10.14306/renhyd.23.3.756

INTRODUCTION

Vegetable fats and oils are widely used as main ingredient in food, medicines, and cosmetics products. In recent years there is a growing demand to exploit newer sources of plant-based oils such as plant seeds that are important sources of oils with high nutritional, industrial and pharmaceutical importance. On the other hand, no oil from a single source has been found to be suitable for all purposes because oils from different sources generally differ in their fatty acid composition (FAC)¹. Nowadays, research on the characterization of by-products in the food processing industry has gained

considerable attention, since the by-products can be utilized in pharmacological, cosmetic, and food applications as potentials source of oils. Fruit seed usually thrown out after processing as an agro-industrial residue, while, they can be used potentially as a natural oil source.

Rambutan (*Nephelium lappaceum* L.), which is a seasonal fruit is believed to originates from Malaysia². It is widely cultivated in Southeast Asian countries such as Thailand (430,000 tons/year), Indonesia (148,000 tons/year), Malaysia (80,000 tons/year), and the Philippines (20,000 tons/year)³. This fruit is generally consumed fresh, but it can be industrially processed to obtain juice, jams, jellies

and marmalades⁴. Rambutan seeds (4-9g/100g) are known as a major waste by product of fruit canning industry⁵. The problems of industrial waste are becoming harder to solve, and much effort will be needed to develop nutritional and industrial potential of by-products, waste, and underutilized agricultural product. It is reported that rambutan seed contains a relatively high amount of fat, with oleic acid and arachidic acid being the major fatty acids^{6,7}. Unlike most vegetable oils and fats, rambutan seed oil contains high level of arachidic acid, which is a long-chain (20 carbons) saturated fatty acid⁶. This characteristic allows the fat to be used without being subjected to hydrogenation and makes it strong resistant to oxidative rancidity⁸. Rambutan seed oil not only could be used for manufacturing candles, soaps, and fuels, but also has a potential to be used in different branch of industries from confectionary products to cosmetics^{7,9}.

Although physicochemical characteristics of rambutan seed fat has been published earlier^{6-8,10-12}, there is no detailed information available for rambutan seed fat varieties grown in the Malaysia. More detailed research on the lipid characteristics of rambutan seed varieties will enhance knowledge and application of rambutan seed oil in a variety of food and specialty products. Thus, the aim of this study was to characterize the physicochemical properties of rambutan seed oil from three rambutan seed varieties (R4, R7 and Serjan) grown in Malaysia to evaluate possible applications of an industrial scale.

MATERIAL AND METHODS

Materials

Three different varieties of fresh and ripe rambutans namely R4, R7 and Serjan were obtained from the University Agricultural Park, Universiti Putra Malaysia (UPM). One batch of samples (each obtained from several trees for each variety during fruiting season which was November 2015–January 2016) was used in this study. The fruits (a total of between 30-35kg for each variety) were manually peeled and deseeded. The seeds represent between 6-8% (~2kg) of the total fruit. Seeds cleaning process was carried out under running tap water and they were then dried in an oven at 60°C for 48 hours. Dried seeds were finely ground using a Waring blender (Model 32BL 80, Dynamic Corporation of America, New Hartford Connecticut, USA) and stored in sealed plastic bags in -20°C for further analysis. All reagents and solvents used were of analytical or HPLC grade

purchased from Merck (Darmstadt, Germany). The standard FAME Mix (C8-C24) was purchased from Supleco (Bellefonte, Pennsylvania, USA).

Oil Extraction

A hundred and fifty grams of ground rambutan seed were placed into a cellulose paper cone and extracted with petroleum ether (b.p 40-60°C) in a 250mL Soxhlet extractor for 8 hours¹³. The solvent was removed via a rotary evaporator Model N-1 (Eyela, Tokyo Rakakikal Co., Ltd., Japan) and residual solvent was removed by drying in an oven at 60°C for 1 hour and flushed with 99.9% nitrogen. The extracted oil was stored at -20°C until it was analysed. Oil extraction was carried out in triplicate for each variety while each analysis was conducted also in triplicate.

Seed Analysis: Moisture content and crude fat content was determined according to AOAC official Methods 930.15 and 945.16, respectively¹³.

Oil Analysis: AOCS Official methods^{14,15} were used to determine free fatty acid (FFA) content (method Ca 5a-40), peroxide value (method Cd 8b-90) and p-anisidine value (method Cd 18-90), iodine value (method Cd 1-25), saponification value (method Cd 3-25), unsaponifiable matter (method Ca 6a-40). PORIM test methods^{16,17} were used to determine viscosity, colour (p4.1) and refractive index (method p4.4). Viscosity was determined at 40°C using Brookfield DV-II viscometer with temperature control (Stoughton, Massachusetts, USA) at 30rpm rotor speed. An oil sample (20mL) was poured into the sample adaptor. The attached spindle of the viscometer was then immersed into the oil. Colour was determined using a Lovibond tintometer Model E (Salisbury, England) according to PORIM test method (p4.1). The oil samples were placed into 1-inch cell and the color was determined at 60°C by achieving the best possible match with the standard colour slides of red and yellow indices. The refractive index (RI) of the oil was determined at 40°C using a digital refractometer Model 300034 (Sper scientific LTD, Scottsdale, Arizona, USA). Frozen rambutan seed oil was first melted to a temperature of 40°C and homogenized. Then, the sample was filtered by a dry Whatman No.1 filter paper to clarify the oil before the determination process. The refractive index was determined at a temperature of 40°C by placing a few drops of the oil in the space between the prisms and determined values were recorded.

Fatty acid composition was determined by conversion of oil to fatty acid methyl esters (FAME) according to the method of Cocks and Van Rede¹⁸ by reacting 50mg of fat with 0.05mL of 1mol/L sodium methoxide in 0.95mL of n-hexane. The

mixtures were vortexed for 5 seconds and allowed to settle for 5 minutes. The top layer (1 μ L) was injected into a gas chromatograph (Model 6890N, Agilent Technologies, Santa Clara, California, USA), equipped with a flame ionization detector (FID) and a polar capillary column BPX70 (0.32mm internal diameter, 30m length and 0.25mm film thickness) (SGE Incorporated, USA) to obtain individual peaks of fatty acid methyl esters. The detector temperature was 250°C and column temperature was 120°C held for 5 minutes and increased at the rate of 8°C/min to 240°C and held for 10 minutes. The run time was 30 minutes. The fatty acid methyl esters peaks were identified by comparing their retention time mixed standard of fatty acids, Supelco FAME mix C8-C24 (Bellefonte, PA, USA). Percent relative fatty acid was calculated based on the peak area of a fatty acid species to the total peak area of all the fatty acids in the oil sample.

A Perkin-Elmer Diamond Differential Scanning Calorimeter (DSC) (Shelton, Connecticut, USA) with PYRIS data processing software was used to determine the thermal properties of oil according to the method used by Abdulkarim *et al.*¹⁹. The instrument was calibrated using indium and zinc. A data processing software, PYRIS Instrument Managing Software, version 10.1, was used to analyse the thermal property of oils. The purge gas was 99.99% nitrogen administered at 100mL/min and 20psi. The oil was first heated in an oven until completely melted. Approximately 3-5mg of oil was placed into a volatile aluminium pan (Perkin Elmer, Shelton, Washington, USA) and hermetically sealed. An empty aluminium pan was used as a reference. Oil samples were subjected to the following programmed temperature: cooling from 60 to -60°C with a cooling rate of 5°C/min and held for 2 minutes; then, heating from -60 to 60°C with a heating rate of 5°C/min and held for another 2 minutes; then, cooling from 60°C to -60°C also at the rate of 5°C/min. The heating and cooling thermograms were recorded and the peak, onset and offset temperature were tabulated. The melting point was determined using melting curve obtained by DSC according to²⁰ Nassu and Gonçalves (1999). The melting

point measured as the temperature at the end of the curve, where the melting phenomenon finishes and all crystals in solid state became liquid.

The solid fat content percentage (SFC%) for all oil samples was derived from the melting thermogram of the respective oil samples according to the method described by Lee *et al.*²¹. The PYRIS data processing software was used to generate the partial peak areas (%) of the melting peaks against the temperature. It is the fact that when the partial area of melting peak was 0%, the SFC of the oil was 100% and vice versa. Thus, by reversing the percentage of partial area from 0-100% to 100-0%, the SFC of the oil sample at different temperature was obtained.

Statistical Analysis

The analytical data were analyzed by one-way analysis of variance followed by Tukey's test using Minitab v. 16 Statistical Software (Minitab Inc., State College, PA, USA). The results were expressed as mean of triplicate determination with standard deviation. Statistical significance differences were considered at the level of $p < 0.05$.

RESULTS

The crude fat content and moisture content of rambutan seed of three varieties are shown in Table 1.

Physicochemical properties such as colour, refractive index (RI), viscosity, melting point, free fatty acid (FFA), peroxide value (PV), p-anisidine value (p-AV), iodine value (IV), saponification value (SV) and unsaponifiable matter (USM) of rambutan seed oil for varieties R4, R7 and Serjan are shown in Table 2.

Table 3 shows the fatty acid composition of rambutan seed oil for varieties R4, R7 and Serjan.

Table 1. Crude fat content and moisture content of rambutan seed for varieties R4, R7 and Serjan.

Parameter	R4	R7	Serjan
Crude fat content (%)	34.25 \pm 0.07 ^c	37.62 \pm 0.10 ^a	35.40 \pm 0.28 ^b
Moisture content (%)	34.07 \pm 0.10 ^a	34.25 \pm 0.07 ^a	34.15 \pm 0.07 ^a

Mean \pm standard deviation.

Means in the same row with different letters are significantly different ($P < 0.05$).

Table 2. Physicochemical properties of rambutan seed oil for varieties R4, R7 and Serjan.

Parameter	R4	R7	Serjan
Lovibond Colour (Red/Yellow)	R: 0.45 ± 0.07 ^a Y: 3.00 ± 0.00 ^a	R: 0.52 ± 0.03 ^a Y: 3.00 ± 0.07 ^a	R: 0.55 ± 0.07 ^a Y: 3.00 ± 0.07 ^a
RI (40°C)	1.4647 ± 0.001 ^a	1.4631 ± 0.001 ^a	1.4620 ± 0.001 ^a
Viscosity (Cp)	49.25 ± 0.35 ^a	50.40 ± 0.84 ^a	50.95 ± 0.77 ^a
Melting point	24.54 ± 0.05 ^c	25.67 ± 0.10 ^b	26.60 ± 0.14 ^a
FFA (as oleic %)	0.62 ± 0.08 ^a	0.56 ± 0.04 ^a	0.68 ± 0.09 ^a
PV (meq O²/kg)	2.00 ± 0.14 ^a	1.82 ± 0.11 ^a	1.74 ± 0.15 ^a
p-AV	0.66 ± 0.02 ^a	0.49 ± 0.08 ^a	0.54 ± 0.06 ^a
IV (g I₂/100g oil)	47.49 ± 0.04 ^a	46.00 ± 0.03 ^b	44.26 ± 0.10 ^c
SV (mg KOH/g oil)	181.75 ± 0.36 ^a	182.10 ± 0.14 ^a	181.69 ± 0.21 ^a
USM (%)	0.67 ± 0.03 ^a	0.82 ± 0.03 ^a	0.72 ± 0.04 ^a

RI: refractive index; **FFA:** free fatty acid; **PV:** peroxide value; **p-AV:** p-anisidine value;
IV: iodine value; **SV:** saponification value; **USM:** unsaponifiable matter.
Mean values with the same letter within the same row are not significantly different (p>0.05).

Table 3. Fatty acid composition of rambutan seed oil for varieties R4, R7 and Serjan.

Fatty acid	Relative percent (%)		
	R4	R7	Serjan
Myristic acid (C14:0)	0.03 ± 0.02 ^a	0.02 ± 0.00 ^a	0.02 ± 0.00 ^a
Palmitic acid (C16:0)	4.74 ± 0.01 ^a	4.58 ± 0.00 ^a	5.00 ± 0.50 ^a
Palmitoleic (C16:1)	0.59 ± 0.00 ^a	0.41 ± 0.00 ^b	0.60 ± 0.00 ^a
Stearic acid (C18:0)	6.35 ± 0.09 ^c	7.99 ± 0.04 ^a	7.30 ± 0.01 ^b
Oleic acid (C18:1)	40.46 ± 0.05 ^a	40.58 ± 0.05 ^a	37.75 ± 0.12 ^b
Linoleic acid (C18:2)	2.31 ± 0.07 ^a	2.01 ± 0.01 ^b	2.08 ± 0.00 ^b
Linolenic acid (C18:3)	0.17 ± 0.00 ^b	0.16 ± 0.00 ^b	0.23 ± 0.00 ^a
Arachidic acid (C20:0)	34.75 ± 0.06 ^b	35.24 ± 0.05 ^b	36.89 ± 0.49 ^a
Gondoic acid (C20:1)	6.69 ± 0.21 ^a	5.78 ± 0.02 ^b	6.04 ± 0.00 ^b
Behenic acid (C22:0)	3.05 ± 0.00 ^a	2.64 ± 0.09 ^b	3.07 ± 0.01 ^a
Erucic acid (C22:1)	0.56 ± 0.01 ^a	0.44 ± 0.02 ^b	0.54 ± 0.01 ^a
Lignoceric acid (C24:0)	0.22 ± 0.12 ^a	0.09 ± 0.01 ^a	0.39 ± 0.09 ^a
SFA (%)	49.16 ± 0.05 ^c	50.56 ± 0.02 ^b	52.68 ± 0.11 ^a
USFA (%)	50.79 ± 0.05 ^a	49.38 ± 0.02 ^b	47.25 ± 0.11 ^c
Ratio of USFA /SFA	1.030 ± 0.00 ^a	0.97 ± 0.00 ^b	0.89 ± 0.00 ^c

SFA: saturated fatty acids; **USFA:** unsaturated fatty acids.
Mean values with the same letter within the same row are not significantly different (p>0.05).

Thermal behavior of rambutan seed oil of varieties R4, R7 and Serjan is shown in Table 4 and Figure 1. There are three peaks in both melting and crystallization curves.

Figure 2 shows solid fat content (%) of rambutan seed oil for varieties R4, R7 and Serjan.

DISCUSSION

Seed analysis (moisture content and crude fat content)

There was no significant difference in the moisture content of the three varieties. The data obtained for three varieties are like values reported by Augustin & Chua²² and Solís-Fuentes *et al.*¹⁰. There were significant differences ($p < 0.05$) in the crude fat content of the seed between three varieties. As it can be seen, among the three varieties R7 has the highest oil content (37.55%) than Serjan (35.6%) and R4 (34.2%) and is like the value reported by Augustin & Chua²². In contrast, in a study by Chai *et al.*⁷, variety R4 had the highest crude fat content than Serjan and R7. The high percentage of the oil in the seed (37.5 %) makes it a distinct potential source for the oil industry. Several authors have reported that the seed yields between 32% and 39% fat^{6-8,10,11,22,23}. Variation can occur in the seed composition and the differences in moisture and fat content reported by different researchers may be from differences in plant variety, ripening stage at harvesting time and cultivation climate and extraction method applied.

Physicochemical properties of rambutan seed oil

There are no significant differences ($p > 0.05$) in colour, refractive index and viscosity for three rambutan varieties of R4, R7 and Serjan. Rambutan seed oil has golden yellow colour. Colour is one of the food properties that it has been widely used as an index of oil quality. Results obtained in this study shows that red colour over yellow colour ratio (R/Y) is less than 0.3 for all three rambutan seed fat indicating that rambutan seed oils appear visually more yellow than red²¹.

The refractive index of rambutan seed oils from varieties R4, R7 and Serjan were 1.4647, 1.4631 and 1.4620, respectively. These values are consistent with the values of Sirisompong *et al.*⁸ and Solís-Fuentes *et al.*¹⁰, who reported that rambutan seed oil had a refractive index of 1.469 and 1.468, respectively. The refractive index of oil is a basic value that is related to molecular weight, fatty acid chain length, degree of conjugation and degree of unsaturation²⁴. High concentrations of the unsaturated fatty acids result in increase of oil refractive index.

Viscosity defines as the resistance of one part of the fluid to move relative to another one. The viscosity of vegetable oils is affected by many of the physical and chemical properties of oils such as the density, molecular weight, melting point and degree of unsaturation²⁵. Generally, viscosity for vegetable oil increases with molecular weight but decreases with increase in unsaturation and temperature²⁶. The viscosity of rambutan seed oils from varieties R4, R7 and Serjan were 49.25, 50.40 and 50.95 at 25°C, respectively.


Table 4. Temperature of phase transition of melting and crystallization of rambutan seed oil.

Variety	Phase transition temperature (°C)				
	T _{Onset}	T ₁	T ₂	T ₃	T _{End}
Melting					
R4	-35.75 ± 0.33 ^a	-32.95 ± 0.36 ^a	5.43 ± 0.06 ^a	23.19 ± 0.11 ^b	24.76 ± 0.20 ^c
R7	-35.63 ± 0.22 ^a	-32.88 ± 0.39 ^a	5.61 ± 0.18 ^a	23.36 ± 0.45 ^b	25.61 ± 0.16 ^b
Serjan	-35.44 ± 0.09 ^a	-33.04 ± 0.04 ^a	5.89 ± 0.09 ^a	25.02 ± 0.09 ^a	26.57 ± 0.07 ^a
Crystallization					
R4	21.19 ± 0.33 ^a	20.51 ± 0.05 ^b	3.00 ± 0.169 ^a	-42.42 ± 0.44 ^a	-45.20 ± 0.94 ^a
R7	21.54 ± 0.22 ^a	20.73 ± 0.33 ^b	3.30 ± 0.12 ^a	-42.69 ± 0.63 ^a	-45.69 ± 0.64 ^a
Serjan	22.79 ± 0.09 ^a	21.75 ± 0.29 ^a	3.70 ± 0.28 ^a	-42.81 ± 0.11 ^a	-45.78 ± 0.24 ^a

Ton: onset temperature; **Tend:** end set temperature.


1, 2, 3, are transition phases, based on Figures 1 (Melting and crystallization curves of rambutan seed oil). Mean values in the same column with same letter are not significantly different ($p > 0.05$)

Figure 1. Melting (a) and Crystallization (b) thermogram of rambutan seed oil of R4, R7 and Serjan.



These results are comparable with the viscosity of some other vegetable oils such as sunflower, corn, soybean, (52,

54.4, 53.4, respectively) which analysed under the same conditions²⁷.

Figure 2. Solid fat content (%) of rambutan seed oil of varieties R4, R7 and Serjan.

Melting point is a significant physical property of fats and oils which is used in identification purpose and many technological applications of fats and oils. According to the results obtained, significant ($P < 0.05$) differences were observed among the three varieties in their melting point (Table 2). Serjan showed a significantly higher melting point (26.60°C) than the R4 and R7 which were 24.54°C and 25.67°C , respectively. This is probably because of presence of high melting TAG in Serjan rather than R4 and R7. These values are within the range of slip melting point (25.1°C to 51.3°C) and (24.8°C and 50.6°C) of various rambutan varieties that were reported by Kheiri and Mohd Som²⁸ and Chai *et al.*⁷. Similar to the findings in this study, Romain *et al.*²⁹ reported that the melting point of rambutan seed fat is 24°C . Sonwai and Ponprachanuvut¹² and Manaf *et al.*⁶ reported that rambutan seed fat has melting point of 39.2°C and 41.55°C , respectively.

Free fatty acid and peroxide value have been frequently used as two of the most important parameters to monitor the quality of edible oils³⁰. Table 2 shows that no significant differences ($P > 0.05$) of free fatty acids was obtained in all three varieties of rambutan seed. According to Kheiri and

Mohd Som²⁸, FFA of rambutan seed fat obtained from 13 rambutan varieties is between 0.32 and 0.52% which are comparable with the results obtained.

The peroxide values for seeds of R4, R7 and Serjan were 2.00, 1.82 and 1.74 meq of O_2/kg of oil, respectively. There are no significant differences ($p > 0.05$) of PV for these seeds. These values were lower than the maximum limit for PV defined by Codex Alimentarius Commission³¹. The commission permits maximum peroxide value of 10 meq of peroxide/kg of oil for vegetable oils. The PV obtained in this study indicates that rambutan seed oil can be stored for a long period without deterioration.

Table 2 shows that there are no significant differences in p-anisidine value for the oils of varieties R4, R7 and Serjan. The p-AV of R4, R7 and Serjan (0.66, 0.49 and 0.54 respectively), is lower than the value (2.00 p-AV) reported by Subramanian *et al.*³², as p-AV of high quality oil is less than 2. Based on the results obtained, it is found that in three varieties the rate of oxidation is very low.

The iodine value indicates the degree of unsaturation of the oil. Iodine value for the R4, R7 and Serjan were significantly

different ($p < 0.05$) (Table 2). The iodine value of R4 (47.49g $I_2/100g$ oil) is significantly ($p < 0.05$) higher than R7 (46.00g $I_2/100g$ oil) and Serjan (44.26g $I_2/100g$ oil) and indicates R4 is high in unsaturated fatty acids. IV is lower in Serjan due to low content of unsaturated fatty acids (47.25%) and high content of arachidic acid (36.89%). The values obtained are in agreements with IV reported by Kheiri and Mohd Som²⁸ for 13 varieties of rambutan ranging from 41.8 to 49.6g $I_2/100g$ oil. The IV obtained are in agreement with those obtained by Manaf *et al.*⁶, Solís-Fuentes *et al.*¹⁰, Lourith *et al.*²³, Mahisanunt *et al.*⁵ and Chai *et al.*⁷ who reported that IV of rambutan seed fat ranged from 44 to 47g $I_2/100g$.

As presented in Table 2, saponification value for three varieties of rambutan seed oil were not significantly different ($p > 0.05$) with values of 181.75, 182.10 and 181.69mg of KOH/g oil for R4, R7 and Serjan, respectively. The SV of these three seeds oil was within the SV range of rambutan seed fat for 13 varieties (157-190mg KOH/g oil) reported by Kheiri and Mohd Som²⁸. Solís-Fuentes *et al.*¹⁰ and Manaf *et al.*⁶ found that SV of rambutan seed fat was 186 and 182.1 mg KOH/g oil, respectively.

The unsaponifiable matter were not significantly different ($p > 0.05$) in the varieties R4, R7 and Serjan (Table 2). The values obtained are within the unsaponifiable matter value range (0.43–0.82%) reported by Kheiri and Mohd Som²⁸ for 13 rambutan varieties. These three varieties have small amount of unsaponifiable matter (0.67–0.82%) which is like value (0.5%) reported for rambutan seed oil by Manaf *et al.*⁶. Sirisompong *et al.*⁸ found that the unsaponifiable of rambutan seed fat was 0.19% which is lower than value obtained in this study probably due to varietal differences.

Twelve fatty acids are identified, where the major fatty acids in three rambutan seeds oil are arachidic acid (34.75–36.89%) and oleic acid (37.75–40.58%). These two fatty acids constitute 74.62–75.82% of the total fatty acids which is consistent with reported data in previous studies^{6–8,10,12,22}. Augustin & Chua²², in a study using three rambutan varieties reported that seeds contained oleic acid and arachidic acid ranging from 37.91–40.15% and 36.14–36.77%, respectively, which is in agreement with the obtained data in this study. Solís-Fuentes *et al.*¹⁰ and Harahap *et al.*¹¹ also reported that rambutan seed fat contains oleic acid 40.3% and 40.45% and arachidic acid 34.5% and 36.36%, respectively which is in accordance with the results reported in this study. The other main fatty acids are palmitic acid, stearic acid and gondoic acid which are present in amounts of 4.47–5%, 6.35–7.99% and 5.78–6.69%, respectively. The oil contains small amounts of myristic acid, palmitoleic acid, linolenic acid, erucic acid and lignoceric acid. Differences in the fatty acid composition

of the seeds may be due to different varieties, different age of maturity of the rambutan fruits and different climate. The total saturated fatty acid of three rambutan seed oils is 49.16–52.68%, which makes it a strong resistant to oxidative rancidity. Since rambutan seed fat is abundant in both oleic acid and arachidic acid, it may also be a good source of nutritious low-calorie fat suitable for human consumption. Studies have shown that oleic acid has blood pressure reducing effect³³, while arachidic acid is less likely to be nutritionally available to the body following hydrolysis by pancreatic lipase due to very high melting temperature (76°C)⁷.

The phase change in DSC relates to different TAG sets due to different fatty acids compositions. The melting and crystallization thermograms of oils from R4, R7 and Serjan seeds showed three distinct endothermic peaks and exothermic peaks (Figure 1), which correspond to a group of high, middle and low melting point triglycerides, respectively^{5,8,10,12}. The onset of crystallization and melting process for three seed oils were R4, 21.19°C and -35.75°C; R7, 21.54°C and -35.63°C; Serjan, 22.79°C and -35.44°C. According to results for melting thermogram, there is a significant difference in endset temperature among R4, R7 and Serjan seeds. The endset temperature shows higher value in Serjan (26.57°C) than R4 (24.76°C) and R7 (25.61°C). This is because of high content of saturated fatty acids in TAG profile of Serjan. Romain *et al.*²⁹ reported that the rambutan seed fat melted completely at below 30°C. However, studies by Manaf *et al.*⁶ and Sonwai and Ponprachanuvut¹² showed that the complete melting point of rambutan seed fat was 40°C and 41.55°C, respectively, while Chai *et al.*⁷ reported that the complete melting temperature of the 11 varieties of rambutan seed fat was between 24.8°C and 50.6°C. Generally, the oils have high content of saturated fatty acids present DSC crystallization and melting profiles at higher temperature regions as compared to the oils have high content of unsaturated fatty acid.

Solid fat content (SFC) is a useful tool to determine the suitability of oils and fats for a particular application³⁴. The SFC profile of rambutan seed oil for varieties R4, R7 and Serjan shows similar trend of thermal properties. It can be seen from the Figure 2 that among three oils, SFC of Serjan and R4 were clearly highest and lowest at all temperature, respectively. The highest content of long chain saturated fatty acids (C20:0) in Serjan could be the reason of its high SFC over the studied temperature range and this inversely apply to R4. The SFC of R4 and R7 seed oil decreased gradually from 0°C to 15°C then decreased more rapidly after 15°C to melt completely at 24.5°C and 25.6°C, respectively (Figure 2). The SFC% of Serjan shows

a gradual decrease from 0°C to 15°C and after 15°C the SFC exhibits sharp melting behaviour until 26.6°C. Based on results, rambutan seed oils are completely liquid at normal ambient temperature because SFC at that temperature was 0%. These results are in good agreement with SFC results of rambutan seed fat found by Romain *et al.*²⁹.

CONCLUSIONS

In this study, three varieties of rambutan seed oil were characterized. The seeds of three rambutan varieties contain relatively high level of crude fat. Rambutan seeds contain a high arachidic acid content which makes the fat very resistant to oxidation. The melting and crystallization curves showed that the oil has three distinct melting and crystallization peaks. The rambutan seed oil melted completely at the range of 24.54°C–26.60°C. Based on these results, modification of rambutan seed fat by blending and interesterification with softer oil or harder fat may lead to products with wider applications. According to the results attained through this study, rambutan seed fat is applicable for a wide variety of industries ranging from confectionery production to cosmetics. Therefore, rambutan seed can be fully utilized and, subsequently, the amount of waste can be minimized.

COMPETING INTERESTS

Authors state that there are no conflicts of interest in preparing the manuscript.

REFERENCES

- (1) Sorkheh K, Kiani S, Sofo A. Wild almond (*Prunus scoparia* L.) as potential oilseed resource for the future: Studies on the variability of its oil content and composition. *Food Chem.* 2016; 212: 58-64.
- (2) Raihana AN, Marikkar J, Amin I, Shuhaimi M. A review on food values of selected tropical fruits' seeds. *Int J Food Prop.* 2015; 18(11): 2380-92.
- (3) Mehdizadeh S, Lasekan O, Muhammad K, Baharin B. Variability in the fermentation index, polyphenols and amino acids of seeds of rambutan (*Nephelium lappaceum* L.) during fermentation. *J Food Compost Anal.* 2015; 37: 128-35.
- (4) Mahmood K, Kamilah H, Alias AK, Ariffin F. Nutritional and therapeutic potentials of rambutan fruit (*Nephelium lappaceum* L.) and the by-products: a review. *J Food Meas Charact.* 2018; 1: 16.
- (5) Mahisanunt B, Jom KN, Matsukawa S, Klinkesorn U. Solvent fractionation of rambutan (*Nephelium lappaceum* L.) kernel fat for production of non-hydrogenated solid fat: Influence of time and solvent type. *J King Saud Univ Sci.* 2017; 29(1): 32-46.
- (6) Manaf YNA, Marikkar JMN, Long K, Ghazali HM. Physico-chemical characterisation of the fat from red-skin rambutan (*Nephelium lappaceum* L.) seed. *J Oleo Sci.* 2013; 62(6): 335-43.
- (7) Chai KF, Mohd Adzahan N, Karim R, Rukayadi Y, Ghazali HM. Characteristics of fat, and saponin and tannin contents of 11 varieties of rambutan (*Nephelium lappaceum* L.) seed. *Int J Food Prop.* 2018; 21(1): 1091-106.
- (8) Sirisompong W, Jirapakkul W, Klinkesorn U. Response surface optimization and characteristics of rambutan (*Nephelium lappaceum* L.) kernel fat by hexane extraction. *LWT-Food Science and Technology.* 2011; 44(9): 1946-51.
- (9) Chai KF, Adzahan NM, Karim R, Rukayadi Y, Ghazali HM. Fat properties and antinutrient content of rambutan (*Nephelium lappaceum* L.) seed during solid-state fermentation of rambutan fruit. *Food Chem.* 2019; 274: 808-15.
- (10) Solís-Fuentes JA, Camey-Ortiz G, Hernández-Medel MdR, Pérez-Mendoza F, Durán-de-Bazúa C. Composition, phase behavior and thermal stability of natural edible fat from rambutan (*Nephelium lappaceum* L.) seed. 2010.
- (11) Harahap SN, Ramli N, Vafaei N, Said M. Physicochemical and nutritional composition of rambutan anak sekolah (*Nephelium lappaceum* L.) seed and seed oil. *Pak J Nutr.* 2012; 11(11): 1073-7.
- (12) Sonwai S, Ponprachanuvut P. Characterization of physicochemical and thermal properties and crystallization behavior of krabok (*Irvingia Malayana*) and rambutan seed fats. *J Oleo Sci.* 2012; 61(12): 671-9.
- (13) AOAC. Official methods of analysis of the Association of Official Analytical Chemists, 14th ed. Washington, DC, 1984.
- (14) Society AOC, Firestone D. Official methods and recommended practices of the American Oil Chemists' Society: AOCS press; 1994.
- (15) AOCS. Official methods and recommended practices of the American Oil Chemists' Society, 6th ed. Champaign, USA: American Oil Chemists' Society; 2009.
- (16) Methods PT. Methods of Test for Palm Oil and Palm Oil Products. 1988.
- (17) PORIM. PORIM test methods. Palm Oil Research Institute of Malaysia. Kuala Lumpur, Malaysia 1995.
- (18) Cocks LV, Rede Cv. Laboratory handbook for oil and fat analysts. Laboratory handbook for oil and fat analysts. 1966.
- (19) Abdulkarim S, Long K, Lai O, Muhammad S, Ghazali H. Some physico-chemical properties of *Moringa oleifera* seed oil extracted using solvent and aqueous enzymatic methods. *Food Chem.* 2005; 93(2): 253-63.
- (20) Nassu RT, Gonçalves LAG. Determination of melting point of vegetable oils and fats by differential scanning calorimetry (DSC) technique. *Grasas y Aceites.* 1999; 50(1): 16-21.

- (21) Lee S, Radu S, Ariffin A, Ghazali H. Physico-chemical characterization of oils extracted from noni, spinach, lady's finger, bitter gourd and mustard seeds, and copra. *Int J Food Prop.* 2015; 18(11): 2508-27.
- (22) Augustin M, Chua B. Composition of rambutan seeds. *Pertanika.* 1988; 11(2): 211-5.
- (23) Lourith N, Kanlayavattanukul M, Mongkonpaibool K, Butsararakool T, Chinmuang T. Rambutan seed as a new promising unconventional source of specialty fat for cosmetics. *Ind Crops Prod.* 2016; 83: 149-54.
- (24) Nehdi IA, Sbihi H, Tan CP, Zarrouk H, Khalil MI, Al-Resayes SI. Characteristics, composition and thermal stability of *Acacia senegal* (L.) Willd. seed oil. *Ind Crops Prod.* 2012; 36(1): 54-8.
- (25) Igwe IO. The effects of temperature on the viscosity of vegetable oils in solution. *Ind Crops Prod.* 2004; 19(2): 185-90.
- (26) Azian MN, Kamal AM, Panau F, Ten W. Viscosity estimation of triacylglycerols and of some vegetable oils, based on their triacylglycerol composition. *Journal of the American Oil Chemists' Society.* 2001; 78(10): 1001-5.
- (27) Santos J, Santos I, Souza A. Effect of heating and cooling on rheological parameters of edible vegetable oils. *Journal of food Engineering.* 2005; 67(4): 401-5.
- (28) Kheiri M, Som MNM. Physico-Chemical Characteristics of Rambutan Kernel Fat. Serdang: MARDI Publication, 1987.
- (29) Romain V, Ngakegni-Limbili AC, Mouloungui Zp, Ouamba J-M. Thermal properties of monoglycerides from *Nephelium lappaceum* L. Oil, as a natural source of saturated and monounsaturated fatty acids. *Ind Eng Chem Res.* 2013; 52(39): 14089-98.
- (30) Nyam K, Tan C, Lai O, Long K, Man YC. Physicochemical properties and bioactive compounds of selected seed oils. *LWT-Food Science and technology.* 2009; 42(8): 1396-403.
- (31) Commission CA. Recommended internal standards edible fats and oils. Edn FAO/WHO, Rome. 1982; 1(11): 1-179.
- (32) Subramanian R, Nandini K, Sheila P, Gopalakrishna A, Raghavarao K, Nakajima M, et al. Membrane processing of used frying oils. *Journal of the American Oil Chemists' Society.* 2000; 77(3): 323.
- (33) Terés S, Barceló-Coblijn G, Benet M, Alvarez R, Bressani R, Halver J, et al. Oleic acid content is responsible for the reduction in blood pressure induced by olive oil. *Proceedings of the National Academy of Sciences.* 2008.
- (34) Sonwai S, Rungprasertphol P, Nantipipat N, Tungvongcharoan S, Laiyangkoon N. Characterization of Coconut Oil Fractions Obtained from Solvent Fractionation Using Acetone. *J Oleo Sci.* 2017; 66(9): 951-61.